

Lmatrix() — Elimination matrix

Description	Syntax	Remarks and examples	Conformability
Diagnostics	Reference	Also see	

Description

`Lmatrix(n)` returns the $n(n+1)/2 \times n^2$ elimination matrix `L` for which $L \cdot \text{vec}(X) = \text{vech}(X)$, where X is an $n \times n$ symmetric matrix.

Syntax

real matrix `Lmatrix(real scalar n)`

Remarks and examples

stata.com

Elimination matrices are frequently used in computing derivatives of functions of symmetric matrices. Section 9.6 of [Lütkepohl \(1996\)](#) lists many useful properties of elimination matrices.

Conformability

`Lmatrix(n)`:
n: 1×1
result: $n(n+1)/2 \times n^2$

Diagnostics

`Lmatrix(n)` aborts with error if *n* is less than 0 or is missing. *n* is interpreted as `trunc(n)`.

Reference

Lütkepohl, H. 1996. *Handbook of Matrices*. New York: Wiley.

Also see

[M-5] [Dmatrix\(\)](#) — Duplication matrix

[M-5] [Kmatrix\(\)](#) — Commutation matrix

[M-5] [vec\(\)](#) — Stack matrix columns

[M-4] [Standard](#) — Functions to create standard matrices

Stata, Stata Press, and Mata are registered trademarks of StataCorp LLC. Stata and Stata Press are registered trademarks with the World Intellectual Property Organization of the United Nations. Other brand and product names are registered trademarks or trademarks of their respective companies. Copyright © 1985–2023 StataCorp LLC, College Station, TX, USA. All rights reserved.

