Title stata.com

spline3() — Cubic spline interpolation

Description Syntax Remarks and examples Conformability
Diagnostics Reference Also see

Description

spline S(x), y returns the coefficients of a cubic natural spline S(x). The elements of x must be strictly monotone increasing.

spline3eval(*spline_info*, x) uses the information returned by spline3() to evaluate and return the spline at the abscissas x. Elements of the returned result are set to missing if outside the range of the spline. x is assumed to be monotonically increasing.

Syntax

```
real matrix spline3(real vector x, real vector y)
real vector spline3eval(real matrix spline_info, real vector x)
```

Remarks and examples

stata.com

spline3() and spline3eval() is a translation into Mata of Herriot and Reinsch (CUBNATSPLINE) (1973).

```
For xx in [x_i, x_{i+1}): S(xx) = \{(d_it + c_i)t + b_i\}t + y_i with t = xx - x_i. spline3() returns (b, c, d, x, y) or, if x and y are row vectors, (b, c, d, x', y').
```

Conformability

```
spline3(x, y):
                         n \times 1
                                           1 \times n
                                     or
                         n \times 1
                                     or
                                           1 \times n
                y:
                         n \times 5
           result:
spline3eval(spline_info, x):
           spline_info:
                                n \times 5
                      x:
                                m \times 1
                                            or
                                                   1 \times m
                  result:
                                m \times 1
                                            or
                                                  1 \times m
```

2

Diagnostics

```
spline3(x, y) requires that x be in ascending order.
```

spline3eval(spline_info, x) requires that x be in ascending order.

Reference

Herriot, J. G., and C. H. Reinsch. 1973. Algorithm 472: Procedures for natural spline interpolation [E1]. *Communications of the ACM* 16: 763–768. https://doi.org/10.1145/362552.362558.

Also see

[M-4] Mathematical — Important mathematical functions

Stata, Stata Press, and Mata are registered trademarks of StataCorp LLC. Stata and Stata Press are registered trademarks with the World Intellectual Property Organization of the United Nations. Other brand and product names are registered trademarks or trademarks of their respective companies. Copyright © 1985–2023 StataCorp LLC, College Station, TX, USA. All rights reserved.

