

Trigonometric functions

[Contents](#)[Functions](#)[References](#)[Also see](#)

Contents

| | |
|--------------------------|--|
| <code>acos(x)</code> | the radian value of the arccosine of x |
| <code>acosh(x)</code> | the inverse hyperbolic cosine of x |
| <code>asin(x)</code> | the radian value of the arcsine of x |
| <code>asinh(x)</code> | the inverse hyperbolic sine of x |
| <code>atan(x)</code> | the radian value of the arctangent of x |
| <code>atan2(y, x)</code> | the radian value of the arctangent of y/x , where the signs of the parameters y and x are used to determine the quadrant of the answer |
| <code>atanh(x)</code> | the inverse hyperbolic tangent of x |
| <code>cos(x)</code> | the cosine of x , where x is in radians |
| <code>cosh(x)</code> | the hyperbolic cosine of x |
| <code>sin(x)</code> | the sine of x , where x is in radians |
| <code>sinh(x)</code> | the hyperbolic sine of x |
| <code>tan(x)</code> | the tangent of x , where x is in radians |
| <code>tanh(x)</code> | the hyperbolic tangent of x |

Functions

`acos(x)`
 Description: the radian value of the arccosine of x
 Domain: -1 to 1
 Range: 0 to π

`acosh(x)`
 Description: the inverse hyperbolic cosine of x

$$\text{acosh}(x) = \ln(x + \sqrt{x^2 - 1})$$

 Domain: 1 to $8.9\text{e}+307$
 Range: 0 to 709.77

`asin(x)`
 Description: the radian value of the arcsine of x
 Domain: -1 to 1
 Range: $-\pi/2$ to $\pi/2$

`asinh(x)`
 Description: the inverse hyperbolic sine of x

$$\text{asinh}(x) = \ln(x + \sqrt{x^2 + 1})$$

 Domain: $-8.9\text{e}+307$ to $8.9\text{e}+307$
 Range: -709.77 to 709.77

2 Trigonometric functions

atan(x)

Description: the radian value of the arctangent of x

Domain: $-8e+307$ to $8e+307$

Range: $-\pi/2$ to $\pi/2$

atan2(y, x)

Description: the radian value of the arctangent of y/x , where the signs of the parameters y and x are used to determine the quadrant of the answer

Domain y : $-8e+307$ to $8e+307$

Domain x : $-8e+307$ to $8e+307$

Range: $-\pi$ to π

atanh(x)

Description: the inverse hyperbolic tangent of x

$$\operatorname{atanh}(x) = \frac{1}{2} \{ \ln(1+x) - \ln(1-x) \}$$

Domain: -1 to 1

Range: $-8e+307$ to $8e+307$

cos(x)

Description: the cosine of x , where x is in radians

Domain: $-1e+18$ to $1e+18$

Range: -1 to 1

cosh(x)

Description: the hyperbolic cosine of x

$$\operatorname{cosh}(x) = \{ \exp(x) + \exp(-x) \} / 2$$

Domain: -709 to 709

Range: 1 to $4.11e+307$

sin(x)

Description: the sine of x , where x is in radians

Domain: $-1e+18$ to $1e+18$

Range: -1 to 1

sinh(x)

Description: the hyperbolic sine of x

$$\operatorname{sinh}(x) = \{ \exp(x) - \exp(-x) \} / 2$$

Domain: -709 to 709

Range: $-4.11e+307$ to $4.11e+307$

tan(x)

Description: the tangent of x , where x is in radians

Domain: $-1e+18$ to $1e+18$

Range: $-1e+17$ to $1e+17$ or *missing*

tanh(x)

Description: the hyperbolic tangent of x

$$\operatorname{tanh}(x) = \{ \exp(x) - \exp(-x) \} / \{ \exp(x) + \exp(-x) \}$$

Domain: $-8e+307$ to $8e+307$

Range: -1 to 1 or *missing*

□ Technical note

The trigonometric functions are defined in terms of *radians*. There are 2π radians in a circle. If you prefer to think in terms of *degrees*, because there are also 360 degrees in a circle, you may convert degrees into radians by using the formula $r = d\pi/180$, where d represents degrees and r represents radians. Stata includes the built-in constant `_pi`, equal to π to machine precision. Thus, to calculate the sine of `theta`, where `theta` is measured in degrees, you could type

```
sin(theta*_pi/180)
```

`atan()` similarly returns radians, not degrees. The arccotangent can be obtained as

```
acot(x) = _pi/2 - atan(x)
```

□

References

Norton, E. C. 2022. [The inverse hyperbolic sine transformation and retransformed marginal effects](#). *Stata Journal* 22: 702–712.

Oldham, K. B., J. C. Myland, and J. Spanier. 2009. *An Atlas of Functions*. 2nd ed. New York: Springer.

Also see

[FN] [Functions by category](#)

[D] [egen](#) — Extensions to generate

[D] [generate](#) — Create or change contents of variable

[M-5] [sin\(\)](#) — Trigonometric and hyperbolic functions

[U] [13.3 Functions](#)

Stata, Stata Press, and Mata are registered trademarks of StataCorp LLC. Stata and Stata Press are registered trademarks with the World Intellectual Property Organization of the United Nations. StataNow and NetCourseNow are trademarks of StataCorp LLC. Other brand and product names are registered trademarks or trademarks of their respective companies. Copyright © 1985–2023 StataCorp LLC, College Station, TX, USA. All rights reserved.



For suggested citations, see the [FAQ on citing Stata documentation](#).