

Mathematical functions

[Contents](#)[Functions](#)[Video example](#)[References](#)[Also see](#)**Contents**

<code>abs(<i>x</i>)</code>	the absolute value of <i>x</i>
<code>ceil(<i>x</i>)</code>	the unique integer <i>n</i> such that $n - 1 < x \leq n$; <i>x</i> (not “.”) if <i>x</i> is missing, meaning that <code>ceil(.a) = .a</code>
<code>cloglog(<i>x</i>)</code>	the complementary log-log of <i>x</i>
<code>comb(<i>n</i>,<i>k</i>)</code>	the combinatorial function $n!/\{k!(n-k)!\}$
<code>digamma(<i>x</i>)</code>	the <code>digamma()</code> function, $d \ln \Gamma(x)/dx$
<code>exp(<i>x</i>)</code>	the exponential function e^x
<code>floor(<i>x</i>)</code>	the unique integer <i>n</i> such that $n \leq x < n + 1$; <i>x</i> (not “.”) if <i>x</i> is missing, meaning that <code>floor(.a) = .a</code>
<code>int(<i>x</i>)</code>	the integer obtained by truncating <i>x</i> toward 0 (thus, <code>int(5.2) = 5</code> and <code>int(-5.8) = -5</code>); <i>x</i> (not “.”) if <i>x</i> is missing, meaning that <code>int(.a) = .a</code>
<code>invcloglog(<i>x</i>)</code>	the inverse of the complementary log-log function of <i>x</i>
<code>invlogit(<i>x</i>)</code>	the inverse of the logit function of <i>x</i>
<code>ln(<i>x</i>)</code>	the natural logarithm, $\ln(x)$
<code>lnfactorial(<i>n</i>)</code>	the natural log of <i>n</i> factorial = $\ln(n!)$
<code>lngamma(<i>x</i>)</code>	$\ln\{\Gamma(x)\}$
<code>log(<i>x</i>)</code>	the natural logarithm, $\ln(x)$; thus, a synonym for <code>ln(<i>x</i>)</code>
<code>log10(<i>x</i>)</code>	the base-10 logarithm of <i>x</i>
<code>logit(<i>x</i>)</code>	the log of the odds ratio of <i>x</i> , $\text{logit}(x) = \ln\{x/(1-x)\}$
<code>max(<i>x</i>₁,<i>x</i>₂,...,<i>x</i>_{<i>n</i>})</code>	the maximum value of <i>x</i> ₁ , <i>x</i> ₂ , ..., <i>x</i> _{<i>n</i>}
<code>min(<i>x</i>₁,<i>x</i>₂,...,<i>x</i>_{<i>n</i>})</code>	the minimum value of <i>x</i> ₁ , <i>x</i> ₂ , ..., <i>x</i> _{<i>n</i>}
<code>mod(<i>x</i>,<i>y</i>)</code>	the modulus of <i>x</i> with respect to <i>y</i>
<code>reldif(<i>x</i>,<i>y</i>)</code>	the “relative” difference $ x - y /(y + 1)$; 0 if both arguments are the same type of extended missing value; <i>missing</i> if only one argument is missing or if the two arguments are two different types of <i>missing</i>
<code>round(<i>x</i>,<i>y</i>)</code> or <code>round(<i>x</i>)</code>	<i>x</i> rounded in units of <i>y</i> or <i>x</i> rounded to the nearest integer if the argument <i>y</i> is omitted; <i>x</i> (not “.”) if <i>x</i> is missing (meaning that <code>round(.a) = .a</code> and that <code>round(.a,<i>y</i>) = .a</code> if <i>y</i> is not missing) and if <i>y</i> is missing, then “.”) is returned
<code>sign(<i>x</i>)</code>	the sign of <i>x</i> : -1 if <i>x</i> < 0, 0 if <i>x</i> = 0, 1 if <i>x</i> > 0, or <i>missing</i> if <i>x</i> is missing
<code>sqrt(<i>x</i>)</code>	the square root of <i>x</i>
<code>sum(<i>x</i>)</code>	the running sum of <i>x</i> , treating missing values as zero
<code>trigamma(<i>x</i>)</code>	the second derivative of <code>lngamma(<i>x</i>)</code> = $d^2 \ln \Gamma(x)/dx^2$
<code>trunc(<i>x</i>)</code>	a synonym for <code>int(<i>x</i>)</code>

Functions

`abs(x)`

Description: the absolute value of x

Domain: $-8e+307$ to $8e+307$

Range: 0 to $8e+307$

`ceil(x)`

Description: the unique integer n such that $n - 1 < x \leq n$; x (not “.”) if x is missing, meaning that `ceil(.a) = .a`

Also see `floor(x)`, `int(x)`, and `round(x)`.

Domain: $-8e+307$ to $8e+307$

Range: integers in $-8e+307$ to $8e+307$

`cloglog(x)`

Description: the complementary log-log of x

$$\text{cloglog}(x) = \ln\{-\ln(1-x)\}$$

Domain: 0 to 1

Range: $-8e+307$ to $8e+307$

`comb(n, k)`

Description: the combinatorial function $n!/\{k!(n-k)!\}$

Domain n : integers 1 to $1e+305$

Domain k : integers 0 to n

Range: 0 to $8e+307$ or *missing*

`digamma(x)`

Description: the `digamma()` function, $d \ln \Gamma(x) / dx$

This is the derivative of `lngamma(x)`. The `digamma(x)` function is sometimes called the psi function, $\psi(x)$.

Domain: $-1e+15$ to $8e+307$

Range: $-8e+307$ to $8e+307$ or *missing*

`exp(x)`

Description: the exponential function e^x

This function is the inverse of `ln(x)`.

Domain: $-8e+307$ to 709

Range: 0 to $8e+307$

`floor(x)`

Description: the unique integer n such that $n \leq x < n + 1$; x (not “.”) if x is missing, meaning that `floor(.a) = .a`

Also see `ceil(x)`, `int(x)`, and `round(x)`.

Domain: $-8e+307$ to $8e+307$

Range: integers in $-8e+307$ to $8e+307$

int(*x*)

Description: the integer obtained by truncating x toward 0 (thus, $\text{int}(5.2) = 5$ and $\text{int}(-5.8) = -5$); x (not “.”) if x is missing, meaning that $\text{int}(.a) = .a$

One way to obtain the closest integer to x is $\text{int}(x + \text{sign}(x)/2)$, which simplifies to $\text{int}(x+0.5)$ for $x \geq 0$. However, use of the `round()` function is preferred. Also see `round(x)`, `ceil(x)`, and `floor(x)`.

Domain: $-8\text{e}+307$ to $8\text{e}+307$

Range: integers in $-8\text{e}+307$ to $8\text{e}+307$

invcloglog(*x*)

Description: the inverse of the complementary log-log function of x

$$\text{invcloglog}(x) = 1 - \exp\{-\exp(x)\}$$

Domain: $-8\text{e}+307$ to $8\text{e}+307$

Range: 0 to 1 or *missing*

invlogit(*x*)

Description: the inverse of the logit function of x

$$\text{invlogit}(x) = \exp(x) / \{1 + \exp(x)\}$$

Domain: $-8\text{e}+307$ to $8\text{e}+307$

Range: 0 to 1 or *missing*

ln(*x*)

Description: the natural logarithm, $\ln(x)$

This function is the inverse of $\exp(x)$. The logarithm of x in base b can be calculated via $\log_b(x) = \log_a(x) / \log_a(b)$. Hence,

$$\log_5(x) = \ln(x) / \ln(5) = \log(x) / \log(5) = \log_{10}(x) / \log_{10}(5)$$

$$\log_2(x) = \ln(x) / \ln(2) = \log(x) / \log(2) = \log_{10}(x) / \log_{10}(2)$$

You can calculate $\log_b(x)$ by using the formula that best suits your needs.

Domain: $1\text{e}-323$ to $8\text{e}+307$

Range: -744 to 709

lnfactorial(*n*)

Description: the natural log of n factorial = $\ln(n!)$

To calculate $n!$, use `round(exp(lnfactorial(n)), 1)` to ensure that the result is an integer. Logs of factorials are generally more useful than the factorials themselves because of overflow problems.

Domain: integers 0 to $1\text{e}+305$

Range: 0 to $8\text{e}+307$

lngamma(*x*)

Description: $\ln\{\Gamma(x)\}$

Here the gamma function, $\Gamma(x)$, is defined by $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$. For integer values of $x > 0$, this is $\ln((x-1)!)$.

`lngamma(x)` for $x < 0$ returns a number such that $\exp(\text{lngamma}(x))$ is equal to the absolute value of the gamma function, $\Gamma(x)$. That is, `lngamma(x)` always returns a real (not complex) result.

Domain: $-2,147,483,648$ to $1\text{e}+305$ (excluding negative integers)

Range: $-8\text{e}+307$ to $8\text{e}+307$

4 Mathematical functions

$\log(x)$

Description: the natural logarithm, $\ln(x)$; thus, a synonym for $\ln(x)$

Domain: $1e-323$ to $8e+307$

Range: -744 to 709

$\log_{10}(x)$

Description: the base-10 logarithm of x

Domain: $1e-323$ to $8e+307$

Range: -323 to 308

$\text{logit}(x)$

Description: the log of the odds ratio of x , $\text{logit}(x) = \ln\{x/(1-x)\}$

Domain: 0 to 1 (exclusive)

Range: $-8e+307$ to $8e+307$ or *missing*

$\max(x_1, x_2, \dots, x_n)$

Description: the maximum value of x_1, x_2, \dots, x_n

Unless all arguments are *missing*, missing values are ignored.

$\max(2, 10, ., 7) = 10$

$\max(., ., .) = .$

Domain x_1 : $-8e+307$ to $8e+307$ or *missing*

Domain x_2 : $-8e+307$ to $8e+307$ or *missing*

...

Domain x_n : $-8e+307$ to $8e+307$ or *missing*

Range: $-8e+307$ to $8e+307$ or *missing*

$\min(x_1, x_2, \dots, x_n)$

Description: the minimum value of x_1, x_2, \dots, x_n

Unless all arguments are *missing*, missing values are ignored.

$\min(2, 10, ., 7) = 2$

$\min(., ., .) = .$

Domain x_1 : $-8e+307$ to $8e+307$ or *missing*

Domain x_2 : $-8e+307$ to $8e+307$ or *missing*

...

Domain x_n : $-8e+307$ to $8e+307$ or *missing*

Range: $-8e+307$ to $8e+307$ or *missing*

$\text{mod}(x, y)$

Description: the modulus of x with respect to y

$\text{mod}(x, y) = x - y \text{ floor}(x/y)$

$\text{mod}(x, 0) = .$

Domain x : $-8e+307$ to $8e+307$

Domain y : 0 to $8e+307$

Range: 0 to $8e+307$

reldif(x, y)

Description: the “relative” difference $|x - y|/(|y| + 1)$; 0 if both arguments are the same type of extended missing value; *missing* if only one argument is missing or if the two arguments are two different types of *missing*

Domain x : $-8e+307$ to $8e+307$ or *missing*

Domain y : $-8e+307$ to $8e+307$ or *missing*

Range: $-8e+307$ to $8e+307$ or *missing*

round(x, y) or **round**(x)

Description: x rounded in units of y or x rounded to the nearest integer if the argument y is omitted; x (not “.”) if x is missing (meaning that $\text{round}(.a) = .a$ and that $\text{round}(.a, y) = .a$ if y is not missing) and if y is missing, then “.” is returned

For $y = 1$, or with y omitted, this amounts to the closest integer to x ; $\text{round}(5.2, 1)$ is 5, as is $\text{round}(4.8, 1)$; $\text{round}(-5.2, 1)$ is -5 , as is $\text{round}(-4.8, 1)$. The rounding definition is generalized for $y \neq 1$. With $y = 0.01$, for instance, x is rounded to two decimal places; $\text{round}(\text{sqrt}(2), .01)$ is 1.41. y may also be larger than 1; $\text{round}(28, 5)$ is 30, which is 28 rounded to the closest multiple of 5. For $y = 0$, the function is defined as returning x unmodified. Also see [int](#)(x), [ceil](#)(x), and [floor](#)(x).

Domain x : $-8e+307$ to $8e+307$

Domain y : $-8e+307$ to $8e+307$

Range: $-8e+307$ to $8e+307$

sign(x)

Description: the sign of x : -1 if $x < 0$, 0 if $x = 0$, 1 if $x > 0$, or *missing* if x is missing

Domain: $-8e+307$ to $8e+307$ or *missing*

Range: $-1, 0, 1$ or *missing*

sqrt(x)

Description: the square root of x

Domain: 0 to $8e+307$

Range: 0 to $1e+154$

sum(x)

Description: the running sum of x , treating missing values as zero

For example, following the command `generate y=sum(x)`, the j th observation on y contains the sum of the first through j th observations on x . See [\[D\] egen](#) for an alternative sum function, `total()`, that produces a constant equal to the overall sum.

Domain: all real numbers or *missing*

Range: $-8e+307$ to $8e+307$ (excluding *missing*)

trigamma(x)

Description: the second derivative of $\text{lngamma}(x) = d^2 \ln \Gamma(x) / dx^2$

The `trigamma()` function is the derivative of `digamma(x)`.

Domain: $-1e+15$ to $8e+307$

Range: 0 to $8e+307$ or *missing*

trunc(x)

Description: a synonym for [int](#)(x)

Video example

[How to round a continuous variable](#)

References

Abramowitz, M., and I. A. Stegun, ed. 1964. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Washington, DC: National Bureau of Standards.

Cox, N. J. 2003. [Stata tip 2: Building with floors and ceilings](#). *Stata Journal* 3: 446–447.

———. 2007. [Stata tip 43: Remainders, selections, sequences, extractions: Uses of the modulus](#). *Stata Journal* 7: 143–145.

Oldham, K. B., J. C. Myland, and J. Spanier. 2009. *An Atlas of Functions*. 2nd ed. New York: Springer.

Also see

[FN] [Functions by category](#)

[D] [egen](#) — Extensions to generate

[D] [generate](#) — Create or change contents of variable

[M-4] [intro](#) — Categorical guide to Mata functions

[U] [13.3 Functions](#)