

rologit — Rank-ordered logistic regression
[Description](#)[Quick start](#)[Menu](#)[Syntax](#)[Options](#)[Remarks and examples](#)[Stored results](#)[Methods and formulas](#)[Acknowledgment](#)[References](#)[Also see](#)

Description

`rologit` fits the rank-ordered logistic regression model by maximum likelihood (Beggs, Cardell, and Hausman 1981). This model is also known as the Plackett–Luce model (Marden 1995), as the exploded logit model (Punj and Staelin 1978), and as the choice-based method of conjoint analysis (Hair et al. 2010).

`rologit` expects the data to be in long form, similar to `clogit` (see [R] `clogit`), in which each of the ranked alternatives forms an observation; all observations related to an individual are linked together by the variable that you specify in the `group()` option. The distinction from `clogit` is that `depvar` in `rologit` records the rankings of the alternatives, whereas for `clogit`, `depvar` marks only the best alternative by a value not equal to zero. `rologit` interprets equal scores of `depvar` as ties. The ranking information may be incomplete “at the bottom” (least preferred alternatives). That is, unranked alternatives may be coded as 0 or as a common value that may be specified with the `incomplete()` option.

If your data record only the unique best alternative, `rologit` fits the same model as `clogit`.

Quick start

Rank-ordered logit model of rankings `y` of alternatives within groups defined by `idvar` using covariates `x1`, `x2`, and `x3`

```
rologit y x1 x2 x3, group(idvar)
```

As above, but interpret the lowest value of `y` as the best

```
rologit y x1 x2 x3, group(idvar) reverse
```

Use Breslow’s method for handling ties in rankings

```
rologit y x1 x2 x3, group(idvar) ties(breslow)
```

With cluster–robust standard errors for clustering by levels of `cvar`

```
rologit y x1 x2 x3, group(idvar) vce(cluster cvar)
```

Menu

Statistics > Ordinal outcomes > Rank-ordered logistic regression

Syntax

```
rologit depvar indepvars [if] [in] [weight], group(varname) [options]
```

<i>options</i>	Description
Model	
* <u>group</u> (<i>varname</i>)	identifier variable that links the alternatives
<u>offset</u> (<i>varname</i>)	include <i>varname</i> in model with coefficient constrained to 1
<u>incomplete</u> (#)	use # to code unranked alternatives; default is <code>incomplete(0)</code>
<u>reverse</u>	reverse the preference order
<u>notest</u> <i>rhs</i>	keep right-hand-side variables that do not vary within group
<u>ties</u> (<i>spec</i>)	method to handle ties: <code>exactm</code> , <code>breslow</code> , <code>efron</code> , or <code>none</code>
SE/Robust	
<u>vce</u> (<i>vcetype</i>)	<i>vcetype</i> may be <code>oim</code> , <code>robust</code> , <code>cluster</code> <i>clustvar</i> , <code>bootstrap</code> , or <code>jackknife</code>
Reporting	
<u>level</u> (#)	set confidence level; default is <code>level(95)</code>
<u>display_options</u>	control columns and column formats, row spacing, line width, display of omitted variables and base and empty cells, and factor-variable labeling
Maximization	
<u>maximize_options</u>	control the maximization process; seldom used
<u>coeflegend</u>	display legend instead of statistics
* <code>group</code> (<i>varname</i>) is required.	
<i>indepvars</i> may contain factor variables; see [U] 11.4.3 Factor variables .	
<code>bootstrap</code> , <code>by</code> , <code>fp</code> , <code>jackknife</code> , <code>rolling</code> , and <code>statsby</code> are allowed; see [U] 11.1.10 Prefix commands .	
Weights are not allowed with the <code>bootstrap</code> prefix; see [R] bootstrap .	
<code>fweights</code> , <code>iwweights</code> , and <code>pweights</code> are allowed, except with <code>ties(efron)</code> ; see [U] 11.1.6 weight .	
<code>coeflegend</code> does not appear in the dialog box.	
See [U] 20 Estimation and postestimation commands for more capabilities of estimation commands.	

Options

Model

`group`(*varname*) is required, and it specifies the identifier variable (numeric or string) that links the alternatives for an individual, which have been compared and rank ordered with respect to one another.

`offset`(*varname*); see [\[R\] estimation options](#).

`incomplete`(#) specifies the numeric value used to code alternatives that are not ranked. It is assumed that unranked alternatives are less preferred than the ranked alternatives (that is, the data record the ranking of the most preferred alternatives). It is not assumed that subjects are indifferent between the unranked alternatives. # defaults to 0.

`reverse` specifies that in the preference order, a higher number means a less attractive alternative. The default is that higher values indicate more attractive alternatives. The rank-ordered logit model

is not symmetric in the sense that reversing the ordering simply leads to a change in the signs of the coefficients.

`notestrhs` suppresses the test that the independent variables vary within (at least some of) the groups. Effects of variables that are always constant are not identified. For instance, a rater's gender cannot directly affect his or her rankings; it could affect the rankings only via an interaction with a variable that does vary over alternatives.

`ties(spec)` specifies the method for handling ties (indifference between alternatives) (see [ST] [stcox](#) for details):

<code>exactm</code>	exact marginal likelihood (default)
<code>breslow</code>	Breslow's method (default if <code>pweights</code> specified)
<code>efron</code>	Efron's method (default if robust VCE)
<code>none</code>	no ties allowed

SE/Robust

`vce(vcetype)` specifies the type of standard error reported, which includes types that are derived from asymptotic theory (`oim`), that are robust to some kinds of misspecification (`robust`), that allow for intragroup correlation (`cluster clustvar`), and that use bootstrap or jackknife methods (`bootstrap`, `jackknife`); see [R] [vce_option](#).

If `ties(exactm)` is specified, `vcetype` may be only `oim`, `bootstrap`, or `jackknife`.

Reporting

`level(#)`; see [R] [estimation options](#).

`display_options:` `noci`, `nopvalues`, `noomitted`, `vsquish`, `noemptycells`, `baselevels`, `allbaselevels`, `nofvlabel`, `fvwrap(#)`, `fvwrapon(style)`, `cformat(%fmt)`, `pformat(%fmt)`, `sformat(%fmt)`, and `nolstretch`; see [R] [estimation options](#).

Maximization

`maximize_options:` `iterate(#)`, `trace`, `[no]log`, `tolerance(#)`, `ltolerance(#)`, `nrtolerance(#)`, and `nonrntolerance`; see [R] [maximize](#). These options are seldom used.

The following option is available with `rologit` but is not shown in the dialog box:

`coeflegend`; see [R] [estimation options](#).

Remarks and examples

[stata.com](http://www.stata.com)

The rank-ordered logit model can be applied to analyze how decision makers combine attributes of alternatives into overall evaluations of the attractiveness of these alternatives. The model generalizes a version of McFadden's choice model without alternative-specific covariates, as fit by the `clogit` command. It uses richer information about the comparison of alternatives, namely, how decision-makers rank the alternatives rather than just specifying the alternative that they like best.

Remarks are presented under the following headings:

- [Examples](#)
- [Comparing respondents](#)
- [Incomplete rankings and ties](#)
- [Clustered choice data](#)
- [Comparison of rologit and clogit](#)
- [On reversals of rankings](#)

Examples

A popular way to study employer preferences for characteristics of employees is the quasiexperimental “vignette method”. As an example, we consider the research by de Wolf on the labor market position of social science graduates (de Wolf 2000). This study addresses how the educational portfolio (for example, general skills versus specific knowledge) affects short-term and long-term labor-market opportunities. De Wolf asked 22 human resource managers (the respondents) to rank order the six most suitable candidates of 20 fictitious applicants and to rank order these six candidates for three jobs, namely, 1) researcher, 2) management trainee, and 3) policy adviser. Applicants were described by 10 attributes, including their age, gender, details of their portfolio, and work experience. In this example, we analyze a subset of the data. Also, to simplify the output, we drop, at random, 10 nonselected applicants per case. The resulting dataset includes 29 cases, consisting of 10 applicants each. The data are in long form: observations correspond to alternatives (the applications), and alternatives that figured in one decision task are identified by the variable `caseid`. We list the observations for `caseid==7`, in which the respondent considered applicants for a social-science research position.

```
. use http://www.stata-press.com/data/r15/evignet
(Vignet study employer prefs (Inge de Wolf 2000))
. list pref female age grades edufit workexp boardexp if caseid==7, noobs
```

pref	female	age	grades	edufit	workexp	boardexp
0	yes	28	A/B	no	none	no
0	no	25	C/D	yes	one year	no
0	no	25	C/D	yes	none	yes
0	yes	25	C/D	no	internship	yes
1	no	25	C/D	yes	one year	yes
2	no	25	A/B	yes	none	no
3	yes	25	A/B	yes	one year	no
4	yes	25	A/B	yes	none	yes
5	no	25	A/B	yes	internship	no
6	yes	28	A/B	yes	one year	yes

Here six applicants were selected. The rankings are stored in the variable `pref`, where a value of 6 corresponds to “best among the candidates”, a value of 5 corresponds to “second-best among the candidates”, etc. The applicants with a ranking of 0 were not among the best six candidates for the job. The respondent was not asked to express his preferences among these four applicants, but by the elicitation procedure, it is known that he ranks these four applicants below the six selected applicants. The best candidate was a female, 28 years old, with education fitting the job, with good grades (A/B), with 1 year of work experience, and with experience being a board member of a fraternity, a sports club, etc. The profiles of the other candidates read similarly. Here the respondent completed the task; that is, he selected and rank ordered the six most suitable applicants. Sometimes the respondent performed only part of the task.

```
. list pref female age grades edufit workexp boardexp if caseid==18, noobs
```

pref	female	age	grades	edufit	workexp	boardexp
0	no	25	C/D	yes	none	yes
0	no	25	C/D	no	internship	yes
0	no	28	C/D	no	internship	yes
0	yes	25	A/B	no	one year	no
2	yes	25	A/B	no	none	yes
2	no	25	A/B	no	none	yes
2	no	25	A/B	no	one year	yes
5	no	25	A/B	no	none	yes
5	no	25	A/B	no	none	yes
5	yes	25	A/B	no	none	no

The respondent selected the six best candidates and segmented these six candidates into two groups: one group with the three best candidates, and a second group of three candidates that were “still acceptable”. The numbers 2 and 5, indicating these two groups, are arbitrary apart from the implied ranking of the groups. The ties between the candidates in a group indicate that the respondent was not able to rank the candidates within the group.

The purpose of the vignette experiment was to explore and test hypotheses about which of the employees’ attributes are valued by employers, how these attributes are weighted depending on the type of job (described by variable `job` in these data), etc. In the psychometric tradition of [Thurstone \(1927\)](#), *value* is assumed to be linear in the attributes, with the coefficients expressing the direction and weight of the attributes. In addition, it is assumed that *valuation* is to some extent a random procedure, captured by an additive random term. For instance, if value depends only on an applicant’s age and gender, we would have

$$\text{value}(\text{female}_i, \text{age}_i) = \beta_1 \text{female}_i + \beta_2 \text{age}_i + \epsilon_i$$

where the random residual, ϵ_i , captures all omitted attributes. Thus $\beta_1 > 0$ means that the employer assigns higher value to a woman than to a man. Given this conceptualization of value, it is straightforward to model the decision (selection) among alternatives or the ranking of alternatives: the alternative with the highest value is selected (chosen), or the alternatives are ranked according to their value. To complete the specification of a model of choice and of ranking, we assume that the random residual ϵ_i follows an “extreme value distribution of type I”, introduced in this context by [Luce \(1959\)](#). This specific assumption is made mostly for computational convenience.

This model is known by many names. Among others, it is known as the rank-ordered logit model in economics ([Beggs, Cardell, and Hausman 1981](#)), as the exploded logit model in marketing research ([Punj and Staelin 1978](#)), as the choice-based conjoint analysis model ([Hair et al. 2010](#)), and as the Plackett–Luce model ([Marden 1995](#)). The model coefficients are estimated using the method of maximum likelihood. The implementation in `rologit` uses an analogy between the rank-ordered logit model and the Cox regression model observed by [Allison and Christakis \(1994\)](#); see [Methods and formulas](#). The `rologit` command implements this method for rankings, whereas `clologit` deals with the variant of choices, that is, only the most highly valued alternative is recorded. In the latter case, the model is also known as the Luce–McFadden choice model. In fact, when the data record the most preferred (unique) alternative and no additional ranking information about preferences is available, `rologit` and `clologit` return the same information, though formatted somewhat differently.

```

. rologit pref female age grades edufit workexp boardexp if job==1, group(caseid)
Iteration 0:  log likelihood =  -95.41087
Iteration 1:  log likelihood =  -71.180903
Iteration 2:  log likelihood =  -68.47734
Iteration 3:  log likelihood =  -68.345918
Iteration 4:  log likelihood =  -68.345389
Refining estimates:
Iteration 0:  log likelihood =  -68.345389

Rank-ordered logistic regression          Number of obs    =    80
Group variable: caseid                  Number of groups =     8
No ties in data                          Obs per group:
                                          min =           10
                                          avg =           10.00
                                          max =           10

                                          LR chi2(6)       =    54.13
                                          Prob > chi2      =    0.0000

Log likelihood = -68.34539

```

pref	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
female	-.4487287	.3671307	-1.22	0.222	-1.168292	.2708343
age	-.0984926	.0820473	-1.20	0.230	-.2593024	.0623172
grades	3.064534	.6148245	4.98	0.000	1.8595	4.269568
edufit	.7658064	.3602366	2.13	0.034	.0597556	1.471857
workexp	1.386427	.292553	4.74	0.000	.8130341	1.959821
boardexp	.6944377	.3762596	1.85	0.065	-.0430176	1.431893

Focusing only on the variables whose coefficients are significant at the 10% level (we are analyzing 8 respondents only!), the estimated value of an applicant for a job of type 1 (research positions) can be written as

$$\text{value} = 3.06 \cdot \text{grades} + 0.77 \cdot \text{edufit} + 1.39 \cdot \text{workexp} + 0.69 \cdot \text{boardexp}$$

Thus employers prefer applicants for a research position ($\text{job}=1$) whose educational portfolio fits the job, who have better grades, who have more relevant work experience, and who have (extracurricular) board experience. They do not seem to care much about the sex and age of applicants, which is comforting.

Given these estimates of the valuation by employers, we consider the probabilities that each of the applications is ranked first. Under the assumption that the ϵ_i are independent and follow an extreme value type I distribution, [Luce \(1959\)](#) showed that the probability, π_i , that alternative i is valued higher than alternatives $2, \dots, k$ can be written in the multinomial logit form

$$\pi_i = \Pr \{ \text{value}_1 > \max(\text{value}_2, \dots, \text{value}_m) \} = \frac{\exp(\text{value}_i)}{\sum_{j=1}^k \exp(\text{value}_j)}$$

The probability of observing a specific ranking can be written as the *product* of such terms, representing a sequential decision interpretation in which the rater first chooses the most preferred alternative, and then the most preferred alternative among the rest, etc.

The probabilities for alternatives to be ranked first are conveniently computed by predict.

```
. predict p if e(sample)
(option pr assumed; conditional probability that alternative is ranked first)
(210 missing values generated)
. sort caseid pref p
. list pref p grades edufit workexp boardexp if caseid==7, noobs
```

pref	p	grades	edufit	workexp	boardexp
0	.0027178	C/D	yes	none	yes
0	.0032275	C/D	no	internship	yes
0	.0064231	A/B	no	none	no
0	.0217202	C/D	yes	one year	no
1	.0434964	C/D	yes	one year	yes
2	.0290762	A/B	yes	none	no
3	.2970933	A/B	yes	one year	no
4	.0371747	A/B	yes	none	yes
5	.1163203	A/B	yes	internship	no
6	.4427504	A/B	yes	one year	yes

There clearly is a positive relation between the stated ranking and the predicted probabilities for alternatives to be ranked first, but the association is not perfect. In fact, we would not have expected a perfect association, as the model specifies a (nondegenerate) probability distribution over the possible rankings of the alternatives. These predictions for sets of 10 candidates can also be used to make predictions for subsets of the alternatives. For instance, suppose that only the last three candidates listed in this table would be available. According to parameter estimates of the rank-ordered logit model, the probability that the last of these candidates is selected equals $0.443 / (0.037 + 0.116 + 0.443) = 0.743$.

Comparing respondents

The `rlogit` model assumes that all respondents, HR managers in large public-sector organizations in The Netherlands, use the *same* valuation function; that is, they apply the same decision weights. This is the substantive interpretation of the assumption that the β 's are constant between the respondents. To probe this assumption, we could test whether the coefficients vary between different groups of respondents. For a metric characteristic of the HR manager, such as `firmsize`, we can consider a trend-model in the valuation weights,

$$\beta_{ij} = \alpha_{i0} + \alpha_{i1}\text{firmsize}_j$$

and we can test that the slopes α_{i1} of `firmsize` are zero.

```

. generate firmsize = employer
. rologit pref edufit grades workexp c.firmsize#c.(edufit grades workexp boardexp)
> if job==1, group(caseid) nolog
Rank-ordered logistic regression      Number of obs      =      80
Group variable: caseid                Number of groups   =       8
No ties in data                       Obs per group:
                                         min =             10
                                         avg  =            10.00
                                         max  =             10
                                         LR chi2(7)       =     57.17
Log likelihood = -66.82346              Prob > chi2        =     0.0000

```

pref	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
edufit	1.29122	1.13764	1.13	0.256	-.9385127	3.520953
grades	6.439776	2.288056	2.81	0.005	1.955267	10.92428
workexp	1.23342	.8065067	1.53	0.126	-.347304	2.814144
c.firmsize# c.edufit	-.0173333	.0711942	-0.24	0.808	-.1568714	.1222048
c.firmsize# c.grades	-.2099279	.1218251	-1.72	0.085	-.4487008	.028845
c.firmsize# c.workexp	.0097508	.0525081	0.19	0.853	-.0931632	.1126649
c.firmsize# c.boardexp	.0382304	.0227545	1.68	0.093	-.0063676	.0828284

```

. testparm c.firmsize#c.(edufit grades workexp boardexp)
( 1) c.firmsize#c.edufit = 0
( 2) c.firmsize#c.grades = 0
( 3) c.firmsize#c.workexp = 0
( 4) c.firmsize#c.boardexp = 0

      chi2( 4) =    7.14
      Prob > chi2 =   0.1288

```

The Wald test that the slopes of the interacted `firmsize` variables are jointly zero provides no evidence upon which we would reject the null hypothesis; that is, we do not find evidence against the assumption of constant valuation weights of the attributes by firms of different size. We did not enter `firmsize` as a predictor variable. Characteristics of the decision-making agent do not vary between alternatives. Thus an additive effect of these characteristics on the valuation of alternatives does *not* affect the agent's ranking of alternatives and his choice. Consequently the coefficient of `firmsize` is not identified. `rologit` would in fact have diagnosed the problem and dropped `firmsize` from the analysis. Diagnosing this problem can slow the estimation considerably; the test may be suppressed by specifying the `notestrhs` option.

Incomplete rankings and ties

`rologit` allows incomplete rankings and ties in the rankings as proposed by Allison and Christakis (1994). `rologit` permits rankings to be incomplete only “at the bottom”; namely, that the ranking of the least attractive alternatives for subjects may not be known—do not confuse this with the situation that a subject is indifferent between these alternatives. This form of incompleteness occurred in the example discussed here, because the respondents were instructed to select and rank

only the top six alternatives. It may also be that respondents refused to rank the alternatives that are very unattractive. `rologit` does not allow other forms of incompleteness, for instance, data in which respondents indicate which of four cars they like best, and which one they like least, but not how they rank the two intermediate cars. Another example of incompleteness that cannot be analyzed with `rologit` is data in which respondents select the three alternatives they like best but are not requested to express their preferences among the three selected alternatives.

`rologit` also permits ties in rankings. `rologit` assumes that if a subject expresses a tie between two or more alternatives, he or she actually holds one particular strict preference ordering, but with all possibilities of a strict ordering consistent with the expressed weak ordering being equally probable. For instance, suppose that a respondent ranks alternative 1 highest. He prefers alternatives 2 and 3 over alternative 4, and he is indifferent between alternatives 2 and 3. We assume that this respondent either has the strict preference ordering $1 > 2 > 3 > 4$ or $1 > 3 > 2 > 4$, with both possibilities being equally likely. From a psychometric perspective, it may actually be more appropriate to also assume that the alternatives 2 and 3 are close; for instance, the difference between the associated valuations (utilities) is less than some threshold or minimally discernible difference. Computationally, however, this is a more demanding model.

Clustered choice data

We have seen that applicants with work experience are in a relatively favorable position. To test whether the effects of work experience vary between the jobs, we can include interactions between the type of job and the attributes of applicants. Such interactions can be obtained using factor variables.

Because some HR managers contributed data for more than one job, we cannot assume that their selection decisions for different jobs are independent. We can account for this by specifying the `vce(cluster clustvar)` option. By treating choice data as incomplete ranking data with only the most preferred alternative marked, `rologit` may be used to estimate the model parameters for clustered choice data.

```

. rologit pref job#c.(female grades edufit workexp), group(caseid)
> vce(cluster employer) nolog
2.job 3.job omitted because of no within-caseid variance

Rank-ordered logistic regression          Number of obs    =       290
Group variable: caseid                   Number of groups =        29
Ties handled via the Efron method        Obs per group:
                                           min =           10
                                           avg =          10.00
                                           max =           10

                                           Wald chi2(12)    =        79.57
                                           Prob > chi2      =         0.0000

Log pseudolikelihood = -296.3855          (Std. Err. adjusted for 22 clusters in employer)

```

pref	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
job						
managemen..	0	(omitted)				
policy ad..	0	(omitted)				
female	-.2286609	.2519883	-0.91	0.364	-.7225489	.2652272
grades	2.812555	.8517878	3.30	0.001	1.143081	4.482028
edufit	.7027757	.2398396	2.93	0.003	.2326987	1.172853
workexp	1.224453	.3396773	3.60	0.000	.5586978	1.890208
job#c.female						
managemen..	.0293815	.4829166	0.06	0.951	-.9171177	.9758808
policy ad..	.1195538	.3688844	0.32	0.746	-.6034463	.8425538
job#c.grades						
managemen..	-2.364247	1.005963	-2.35	0.019	-4.335898	-.3925961
policy ad..	-1.88232	.8995277	-2.09	0.036	-3.645362	-.1192782
job#c.edufit						
managemen..	-.267475	.4244964	-0.63	0.529	-1.099473	.5645226
policy ad..	-.3182995	.3689972	-0.86	0.388	-1.041521	.4049217
job#						
c.workexp						
managemen..	-.6870077	.3692946	-1.86	0.063	-1.410812	.0367964
policy ad..	-.4656993	.4515712	-1.03	0.302	-1.350763	.4193639

The parameter estimates for the first job type are very similar to those that would have been obtained from an analysis isolated to these data. Differences are due only to an implied change in the method of handling ties. With clustered observations, `rologit` uses Efron's method. If we had specified the `ties(efron)` option with the separate analyses, then the parameter estimates would have been identical to the simultaneous results. Another difference is that `rologit` now reports robust standard errors, adjusted for clustering within respondents. These could have been obtained for the separate analyses, as well by specifying the `vce(robust)` option. In fact, this option would also have forced `rologit` to switch to Efron's method as well.

Given the combined results for the three types of jobs, we can test easily whether the weights for the attributes of applicants vary between the jobs, in other words, whether employers are looking for different qualifications in applicants for different jobs. A Wald test for the equality hypothesis of no difference can be obtained with the `testparm` command:

```
. testparm job#c.(female grades edufit workexp)
( 1) 2.job#c.female = 0
( 2) 3.job#c.female = 0
( 3) 2.job#c.grades = 0
( 4) 3.job#c.grades = 0
( 5) 2.job#c.edufit = 0
( 6) 3.job#c.edufit = 0
( 7) 2.job#c.workexp = 0
( 8) 3.job#c.workexp = 0

      chi2( 8) =    14.96
      Prob > chi2 =    0.0599
```

We find only mild evidence that employers look for different qualities in candidates according to the job for which they are being considered.

□ Technical note

Allison (1999) stressed that the comparison between groups of the coefficients of logistic regression is problematic, especially in its latent-variable interpretation. In many common latent-variable models, only the regression coefficients divided by the scale of the latent variable are identified. Thus a comparison of logit regression coefficients between, say, men and women is meaningful only if one is willing to argue that the standard deviation of the latent residual does not differ between the sexes. The rank-ordered logit model is also affected by this problem. While we formulated the model with a scale-free residual, we can actually think of the model for the value of an alternative as being scaled by the standard deviation of the random term, representing other relevant attributes of alternatives. Again comparing attribute weights between jobs is meaningful to the extent that we are willing to defend the proposition that “all omitted attributes” are equally important for different kinds of jobs. □

Comparison of rologit and clogit

The rank-ordered logit model also has a sequential interpretation. A subject first chooses the best among the alternatives. Next he or she selects the best alternative among the remaining alternatives, etc. The decisions at each of the subsequent stages are described by a conditional logit model, and a subject is assumed to apply the same decision weights at each stage. Some authors have expressed concern that later choices may well be made more randomly than the first few decisions. A formalization of this idea is a heteroskedastic version of the rank-ordered logit model in which the scale of the random term increases with the number of decisions made (for example, Hausman and Ruud [1987]). This extended model is currently not supported by `rologit`. However, the hypothesis that the same decision weights are applied at the first stage and at later stages can be tested by applying a Hausman test.

First, we fit the rank-ordered logit model on the full ranking data for the first type of job,

```
. rologit pref age female edufit grades workexp boardexp if job==1,
> group(caseid) nolog
```

Rank-ordered logistic regression	Number of obs	=	80
Group variable: caseid	Number of groups	=	8
No ties in data	Obs per group:		
	min	=	10
	avg	=	10.00
	max	=	10
	LR chi2(6)	=	54.13
Log likelihood = -68.34539	Prob > chi2	=	0.0000

pref	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
age	-.0984926	.0820473	-1.20	0.230	-.2593024 .0623172
female	-.4487287	.3671307	-1.22	0.222	-1.168292 .2708343
edufit	.7658064	.3602366	2.13	0.034	.0597556 1.471857
grades	3.064534	.6148245	4.98	0.000	1.8595 4.269568
workexp	1.386427	.292553	4.74	0.000	.8130341 1.959821
boardexp	.6944377	.3762596	1.85	0.065	-.0430176 1.431893

and we save the estimates for later use with the `estimates` command.

```
. estimates store Ranking
```

To estimate the decision weights on the basis of the most preferred alternatives only, we create a variable, `best`, that is 1 for the best alternatives, and 0 otherwise. The `by` prefix is useful here.

```
. by caseid (pref), sort: generate best = pref == pref[_N] if job==1
(210 missing values generated)
```

By specifying `(pref)` with `by caseid`, we ensured that the data were sorted in increasing order on `pref` within `caseid`. Hence, the most preferred alternatives are last in the sort order. The expression `pref == pref[_N]` is true (1) for the most preferred alternatives, even if the alternative is not unique, and false (0) otherwise. If the most preferred alternatives were sometimes tied, we could still fit the model for the based-alternatives-only data via `rologit`, but `clogit` would yield different results because it deals with ties in a less appropriate way for continuous valuations. To ascertain whether there are ties in the selected data regarding applicants for research positions, we can combine `by` with `assert`:

```
. by caseid (pref), sort: assert pref[_N-1] != pref[_N] if job==1
```

There are no ties. We can now fit the model on the choice data by using either `clogit` or `rologit`.

```
. rologit best age edufit grades workexp boardexp if job==1, group(caseid) nolog
Rank-ordered logistic regression          Number of obs   =          80
Group variable: caseid                   Number of groups =           8
No ties in data                          Obs per group:
                                          min =          10
                                          avg =         10.00
                                          max =          10
                                          LR chi2(5)      =         17.27
                                          Prob > chi2     =         0.0040
Log likelihood = -9.783205
```

best	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
age	-.1048959	.2017068	-0.52	0.603	-.5002339	.2904421
edufit	.4558387	.9336775	0.49	0.625	-1.374136	2.285813
grades	3.443851	1.969002	1.75	0.080	-.4153223	7.303025
workexp	2.545648	1.099513	2.32	0.021	.3906422	4.700655
boardexp	1.765176	1.112763	1.59	0.113	-.4157988	3.946152

```
. estimates store Choice
```

The same results, though with a slightly different formatted header, would have been obtained by using `clgfit` on these data.

```
. clgfit best age edufit grades workexp boardexp if job==1, group(caseid) nolog
Conditional (fixed-effects) logistic regression
                                          Number of obs   =          80
                                          LR chi2(5)      =         17.27
                                          Prob > chi2     =         0.0040
Log likelihood = -9.7832046              Pseudo R2       =         0.4689
```

best	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
age	-.1048959	.2017068	-0.52	0.603	-.5002339	.2904421
edufit	.4558387	.9336775	0.49	0.625	-1.374136	2.285813
grades	3.443851	1.969002	1.75	0.080	-.4153223	7.303025
workexp	2.545648	1.099513	2.32	0.021	.3906422	4.700655
boardexp	1.765176	1.112763	1.59	0.113	-.4157988	3.946152

The parameters of the ranking and choice models look different, but the standard errors based on the choice data are much larger. Are we estimating parameters with the ranking data that are different from those with the choice data? A Hausman test compares two estimators of a parameter. One of the estimators should be efficient under the null hypothesis, namely, that choosing the second-best alternative is determined with the same decision weights as the best, etc. In our case, the efficient estimator of the decision weights uses the ranking information. The other estimator should be consistent, even if the null hypothesis is false. In our application, this is the estimator that uses the first-choice data only.

```
. hausman Choice Ranking
```

	Coefficients		(b-B) Difference	sqrt(diag(V_b-V_B)) S.E.
	(b) Choice	(B) Ranking		
age	-.1048959	-.0984926	-.0064033	.1842657
edufit	.4558387	.7658064	-.3099676	.8613846
grades	3.443851	3.064534	.3793169	1.870551
workexp	2.545648	1.386427	1.159221	1.059878
boardexp	1.765176	.6944377	1.070739	1.04722

```

      b = consistent under Ho and Ha; obtained from rologit
      B = inconsistent under Ha, efficient under Ho; obtained from rologit
Test: Ho: difference in coefficients not systematic
      chi2(5) = (b-B)'[(V_b-V_B)^(-1)](b-B)
              = 3.05
      Prob>chi2 = 0.6918

```

We do not find evidence for misspecification. We have to be cautious, though, because Hausman-type tests are often not powerful, and the number of observations in our example is very small, which makes the quality of the method of the null distribution by a chi-squared test rather uncertain.

On reversals of rankings

The rank-ordered logit model has a property that you may find unexpected and even unfortunate. Compare two analyses with the rank-ordered logit model, one in which alternatives are ranked from “most attractive” to “least attractive”, the other a reversed analysis in which these alternatives are ranked from “most unattractive” to “least unattractive”. By unattractiveness, you probably mean just the opposite of attractiveness, and you expect that the weights of the attributes in predicting “attractiveness” to be minus the weights in predicting “unattractiveness”. This is, however, *not* true for the rank-ordered logit model. The assumed distribution of the random residual takes the form $F(\epsilon) = 1 - \exp\{\exp(-\epsilon)\}$. This distribution is right-skewed. Therefore, slightly different models result from adding and subtracting the random residual, corresponding with high-to-low and low-to-high rankings. Thus the estimated coefficients will differ between the two specifications, though usually not in an important way. You may observe the difference by specifying the `reverse` option of `rologit`. Reversing the rank order makes rankings that are incomplete at the bottom become incomplete at the top. Only the first kind of incompleteness is supported by `rologit`. Thus, for this comparison, we exclude the alternatives that are not ranked, omitting the information that ranked alternatives are preferred over excluded ones.

```

. rologit pref grades edufit workexp boardexp if job==1 & pref!=0, group(caseid)
  (output omitted)
. estimates store Original
. rologit pref grades edufit workexp boardexp if job==1 & pref!=0, group(caseid)
> reverse
  (output omitted)
. estimates store Reversed

```

```
. estimates table Original Reversed, stats(aic bic)
```

Variable	Original	Reversed
grades	2.0032332	-1.0955335
edufit	-.13111006	-.05710681
workexp	1.2805373	-1.2096383
boardexp	.46213212	-.27200317
aic	96.750452	99.665642
bic	104.23526	107.15045

Thus, although the weights of the attributes for reversed rankings are indeed mostly of opposite signs, the magnitudes of the weights and their standard errors differ. Which one is more appropriate? We have no advice to offer here. The specific science of the problem will determine what is appropriate, though we would be surprised indeed if this helps here. Formal testing does not help much either, as the models for the original and reversed rankings are not nested. The model-selection indices, such as the AIC and BIC, however, suggest that you stick to the rank-ordered logit model applied to the original ranking rather than to the reversed ranking.

Stored results

`rologit` stores the following in `e()`:

Scalars

<code>e(N)</code>	number of observations
<code>e(ll_0)</code>	log likelihood of the null model (“all rankings are equiprobable”)
<code>e(ll)</code>	log likelihood
<code>e(df_m)</code>	model degrees of freedom
<code>e(chi2)</code>	χ^2
<code>e(p)</code>	significance
<code>e(r2_p)</code>	pseudo- R^2
<code>e(N_g)</code>	number of groups
<code>e(g_min)</code>	minimum group size
<code>e(g_avg)</code>	average group size
<code>e(g_max)</code>	maximum group size
<code>e(code_inc)</code>	value for incomplete preferences
<code>e(N_clust)</code>	number of clusters
<code>e(rank)</code>	rank of <code>e(V)</code>

Macros

<code>e(cmd)</code>	<code>rologit</code>
<code>e(cmdline)</code>	command as typed
<code>e(depvar)</code>	name of dependent variable
<code>e(group)</code>	name of <code>group()</code> variable
<code>e(wtype)</code>	weight type
<code>e(wexp)</code>	weight expression
<code>e(title)</code>	title in estimation output
<code>e(clustvar)</code>	name of cluster variable
<code>e(offset)</code>	linear offset variable
<code>e(chi2type)</code>	Wald or LR; type of model χ^2 test
<code>e(reverse)</code>	<code>reverse</code> , if specified
<code>e(ties)</code>	<code>breslow</code> , <code>efron</code> , <code>exactm</code>
<code>e(vce)</code>	<code>vcetype</code> specified in <code>vce()</code>
<code>e(vcetype)</code>	title used to label Std. Err.
<code>e(properties)</code>	<code>b V</code>
<code>e(predict)</code>	program used to implement <code>predict</code>
<code>e(marginsok)</code>	predictions allowed by margins
<code>e(marginsnotok)</code>	predictions disallowed by margins

<code>e(marginsdefault)</code>	default <code>predict()</code> specification for margins
<code>e(asbalanced)</code>	factor variables <code>fvset</code> as <code>asbalanced</code>
<code>e(asobserved)</code>	factor variables <code>fvset</code> as <code>asobserved</code>

Matrices

<code>e(b)</code>	coefficient vector
<code>e(V)</code>	variance–covariance matrix of the estimators
<code>e(V_modelbased)</code>	model-based variance

Functions

<code>e(sample)</code>	marks estimation sample
------------------------	-------------------------

Methods and formulas

Allison and Christakis (1994) demonstrate that maximum likelihood estimates for the rank-ordered logit model can be obtained as the maximum partial-likelihood estimates of an appropriately specified Cox regression model for waiting time ([ST] `stcox`). In this analogy, a higher value for an alternative is formally equivalent to a higher hazard rate of failure. `rologit` uses `stcox` to fit the rank-ordered logit model based on such a specification of the data in Cox terms. A higher stated preference is represented by a shorter waiting time until failure. Incomplete rankings are dealt with via censoring. Moreover, decision situations (subjects) are to be treated as strata. Finally, as proposed by Allison and Christakis, ties in rankings are handled by the marginal-likelihood method, specifying that all strict preference orderings consistent with the stated weak preference ordering are equally likely. The marginal-likelihood estimator is available in `stcox` via the `exactm` option. The methods of the marginal likelihood due to Breslow and Efron are also appropriate for the analysis of rank-ordered logit models. Because in most applications the number of ranked alternatives by one subject will be fairly small (at most, say, 20), the number of ties is small as well, and so you rarely will need to turn to methods to restrict computer time. Because the marginal-likelihood estimator in `stcox` does not support the cluster adjustment or `pweights`, you should use the Efron method in such cases.

This command supports the clustered version of the Huber/White/sandwich estimator of the variance using `vce(robust)` and `vce(cluster clustvar)`. See [P] `_robust`, particularly *Maximum likelihood estimators* and *Methods and formulas*. Specifying `vce(robust)` is equivalent to specifying `vce(cluster groupvar)`, where *groupvar* is the identifier variable that links the alternatives.

Acknowledgment

The `rologit` command was written by Jeroen Weesie of the Department of Sociology at Utrecht University, The Netherlands.

References

- Allison, P. D. 1999. Comparing logit and probit coefficients across groups. *Sociological Methods and Research* 28: 186–208.
- Allison, P. D., and N. Christakis. 1994. Logit models for sets of ranked items. In Vol. 24 of *Sociological Methodology*, ed. P. V. Marsden, 123–126. Oxford: Blackwell.
- Beggs, S., S. Cardell, and J. A. Hausman. 1981. Assessing the potential demand for electric cars. *Journal of Econometrics* 17: 1–19.
- de Wolf, I. 2000. *Opleidingspecialisatie en arbeidsmarktsucces van sociale wetenschappers*. Amsterdam: ThelaThesis.
- Hair, J. F., Jr., W. C. Black, B. J. Babin, and R. E. Anderson. 2010. *Multivariate Data Analysis*. 7th ed. Upper Saddle River, NJ: Pearson.

- Hausman, J. A., and P. A. Ruud. 1987. Specifying and testing econometric models for rank-ordered data. *Journal of Econometrics* 34: 83–104.
- Luce, R. D. 1959. *Individual Choice Behavior: A Theoretical Analysis*. New York: Dover.
- Marden, J. I. 1995. *Analyzing and Modeling Rank Data*. London: Chapman & Hall.
- McCullagh, P. 1993. Permutations and regression models. In *Probability Models and Statistical Analysis for Ranking Data*, ed. M. A. Fligner and J. S. Verducci, 196–215. New York: Springer.
- Plackett, R. L. 1975. The analysis of permutations. *Applied Statistics* 24: 193–202.
- Punj, G. N., and R. Staelin. 1978. The choice process for graduate business schools. *Journal of Marketing Research* 15: 588–598.
- Thurstone, L. L. 1927. A law of comparative judgment. *Psychological Reviews* 34: 273–286.
- Yellott, J. I., Jr. 1977. The relationship between Luce’s choice axiom, Thurstone’s theory of comparative judgment, and the double exponential distribution. *Journal of Mathematical Psychology* 15: 109–144.

Also see

- [R] **rologit postestimation** — Postestimation tools for rologit
- [R] **clomit** — Conditional (fixed-effects) logistic regression
- [R] **logistic** — Logistic regression, reporting odds ratios
- [R] **mlogit** — Multinomial (polytomous) logistic regression
- [R] **nlogit** — Nested logit regression
- [R] **slogit** — Stereotype logistic regression
- [U] **20 Estimation and postestimation commands**