

**zttest** —  $z$  tests (mean-comparison tests, known variance)

<a href="#">Description</a>	<a href="#">Quick start</a>	<a href="#">Menu</a>	<a href="#">Syntax</a>
<a href="#">Options</a>	<a href="#">Remarks and examples</a>	<a href="#">Stored results</a>	<a href="#">Methods and formulas</a>
<a href="#">References</a>	<a href="#">Also see</a>		

## Description

`zttest` performs  $z$  tests on the equality of means, assuming known variances. The test can be performed for one sample against a hypothesized population value or for no difference in population means estimated from two samples. Two-sample tests can be conducted for paired and unpaired data. Clustered data are also supported.

`zttesti` is the immediate form of `zttest`; see [\[U\] 19 Immediate commands](#).

For the comparison of means when variances are unknown, use `ttest`; see [\[R\] ttest](#).

## Quick start

One-sample test that the mean of `v1` is 3 at the 90% confidence level

```
zttest v1 == 3, level(90)
```

As above, and adjust for clustering with clusters defined by `cvar` and an intraclass correlation of 0.5

```
zttest v1 == 3, level(90) cluster(cvar) rho(0.5)
```

Unpaired  $z$  test that the mean of `v1` is equal between two groups defined by `catvar`

```
zttest v1, by(catvar)
```

As above, and adjust for clustering with clusters defined by `cvar` and an intraclass correlation of 0.5 in the two groups

```
zttest v1, by(catvar) cluster(cvar) rho(0.5)
```

Unpaired test of equality of the means of `v2` and `v3`

```
zttest v2 == v3, unpaired
```

Paired test of equality of the means of `v2` and `v3` with standard deviation of the differences between paired observations of 2.4

```
zttest v2 == v3, sddiff(2.4)
```

As above, specified using a common standard deviation of 2 and correlation between observations of 0.28

```
zttest v2 == v3, sd(2) corr(0.28)
```

Immediate form unpaired test of  $\mu_1 = \mu_2$  if  $\bar{x}_1 = 3.2$ ,  $sd_1 = 0.1$ ,  $\bar{x}_2 = 3.4$ , and  $sd_2 = 0.15$  with  $n_1 = n_2 = 120$

```
zttesti 120 3.2 0.1 120 3.4 0.15
```

# Menu

## ztest

Statistics > Summaries, tables, and tests > Classical tests of hypotheses > z test (mean-comparison test, known variance)

## ztesti

Statistics > Summaries, tables, and tests > Classical tests of hypotheses > z test calculator

# Syntax

### One-sample z test

```
ztest varname == # [if] [in] [, onesampleopts]
```

### Two-sample z test using groups

```
ztest varname [if] [in], by(groupvar) [twosamplegropts]
```

### Two-sample z test using variables

```
ztest varname1 == varname2 [if] [in], unpaired [twosamplevaropts]
```

### Paired z test

```
ztest varname1 == varname2 [if] [in], sddiff(#) [level(#)]
```

```
ztest varname1 == varname2 [if] [in], corr(#) [pairedopts]
```

### Immediate form of one-sample z test

```
ztesti #obs #mean #sd #val [, level(#)]
```

### Immediate form of two-sample unpaired z test

```
ztesti #obs1 #mean1 #sd1 #obs2 #mean2 #sd2 [, level(#)]
```

### onesampleopts

### Description

---

#### Main

sd(#)	one-population standard deviation; default is sd(1)
level(#)	confidence level; default is level(95)
cluster(varname)	variable defining the clusters
rho(#)	intraclass correlation

---

---

*twosamplegopts* Description

---

Main

* <i>by</i> ( <i>groupvar</i> )	variable defining the groups
<u>unpaired</u>	unpaired test; implied when <i>by</i> () is specified
<i>sd</i> (#)	two-population common standard deviation; default is <i>sd</i> (1)
<i>sd1</i> (#)	standard deviation of the first population; requires <i>sd2</i> () and may not be combined with <i>sd</i> ()
<i>sd2</i> (#)	standard deviation of the second population; requires <i>sd1</i> () and may not be combined with <i>sd</i> ()
<u>level</u> (#)	confidence level; default is <i>level</i> (95)
<i>cluster</i> ( <i>varname</i> )	variable defining the clusters
<i>rho</i> (#)	common intraclass correlation
<i>rho1</i> (#)	intraclass correlation for group 1
<i>rho2</i> (#)	intraclass correlation for group 2

---

\**by*(*groupvar*) is required.

---

*twosamplevaropts* Description

---

Main

* <u>unpaired</u>	unpaired test
<i>sd</i> (#)	two-population common standard deviation; default is <i>sd</i> (1)
<i>sd1</i> (#)	standard deviation of the first population; requires <i>sd2</i> () and may not be combined with <i>sd</i> ()
<i>sd2</i> (#)	standard deviation of the second population; requires <i>sd1</i> () and may not be combined with <i>sd</i> ()
<u>level</u> (#)	confidence level; default is <i>level</i> (95)

---

\*unpaired is required.

---

*pairedopts* Description

---

Main

* <i>corr</i> (#)	correlation between paired observations
<i>sd</i> (#)	two-population common standard deviation; default is <i>sd</i> (1); may not be combined with <i>sd1</i> (), <i>sd2</i> (), or <i>sddiff</i> ()
<i>sd1</i> (#)	standard deviation of the first population; requires <i>corr</i> () and <i>sd2</i> () and may not be combined with <i>sd</i> () or <i>sddiff</i> ()
<i>sd2</i> (#)	standard deviation of the second population; requires <i>corr</i> () and <i>sd1</i> () and may not be combined with <i>sd</i> () or <i>sddiff</i> ()
<u>level</u> (#)	confidence level; default is <i>level</i> (95)

---

\**corr*(#) is required.*by* is allowed with *ztest*; see [D] *by*.

## Options

Main

`by(groupvar)` specifies the *groupvar* that defines the two groups that `ztest` will use to test the hypothesis that their means are equal. Specifying `by(groupvar)` implies an unpaired (two-sample) *z* test. Do not confuse the `by()` option with the `by` prefix; you can specify both.

`unpaired` specifies that the data be treated as unpaired. The `unpaired` option is used when the two sets of values to be compared are in different variables.

`sddiff(#)` specifies the population standard deviation of the differences between paired observations for a paired *z* test. For this kind of test, either `sddiff()` or `corr()` must be specified.

`corr(#)` specifies the correlation between paired observations for a paired *z* test. This option along with `sd1()` and `sd2()` or with `sd()` is used to compute the standard deviation of the differences between paired observations unless that standard deviation is supplied directly in the `sddiff()` option. For a paired *z* test, either `sddiff()` or `corr()` must be specified.

`sd(#)` specifies the population standard deviation for a one-sample *z* test or the common population standard deviation for a two-sample *z* test. The default is `sd(1)`. `sd()` may not be combined with `sd1()`, `sd2()`, or `sddiff()`.

`sd1(#)` specifies the standard deviation of the first population or group. When `sd1()` is specified with `by(groupvar)`, the first group is defined by the first category of the sorted *groupvar*. `sd1()` requires `sd2()` and may not be combined with `sd()` or `sddiff()`.

`sd2(#)` specifies the standard deviation of the second population or group. When `sd2()` is specified with `by(groupvar)`, the second group is defined by the second category of the sorted *groupvar*. `sd2()` requires `sd1()` and may not be combined with `sd()` or `sddiff()`.

`level(#)` specifies the confidence level, as a percentage, for confidence intervals. The default is `level(95)` or as set by `set level`; see [U] 20.8 **Specifying the width of confidence intervals**.

`cluster(varname)` specifies the variable that identifies clusters. The `cluster()` option is required to adjust the computation for clustering.

`rho(#)` specifies the intraclass correlation for a one-sample test or the common intraclass correlation for a two-sample test. The `rho()` option is required to adjust the computation for clustering for a one-sample test.

`rho1(#)` specifies the intraclass correlation of the first group for a two-sample test using groups. The `rho()` option or both `rho1()` and `rho2()` options are required to adjust the computation for clustering.

`rho2(#)` specifies the intraclass correlation of the second group for a two-sample test using groups. The `rho()` option or both `rho1()` and `rho2()` options are required to adjust the computation for clustering.

When `by()` is used, `sd1()` and `sd2()` or `sd()` is used to specify the population standard deviations of the two groups defined by *groupvar* for an unpaired two-sample *z* test (using groups). By default, a common standard deviation of one, `sd(1)`, is assumed.

When `unpaired` is used, `sd1()` and `sd2()` or `sd()` is used to specify the population standard deviations of *varname*<sub>1</sub> and *varname*<sub>2</sub> for an unpaired two-sample *z* test (using variables). By default, a common standard deviation of one, `sd(1)`, is assumed.

Options `corr()`, `sd1()`, and `sd2()` or `corr()` and `sd()` are used for a paired *z* test to compute the standard deviation of the differences between paired observations. By default, a common standard

deviation of one, `sd(1)`, is assumed for both populations. Alternatively, the standard deviation of the differences between paired observations may be supplied directly with the `sddiff()` option.

## Remarks and examples

[stata.com](http://www.stata.com)

Remarks are presented under the following headings:

*One-sample z test*  
*Two-sample z test*  
*Paired z test*  
*Adjust for clustering*  
*Immediate form*

For the purpose of illustration, we assume that variances are known in all the examples below.

## One-sample z test

### ► Example 1

In the first form, `ztest` tests whether the mean of the sample is equal to a known constant under the assumption of known variance. Assume that we have a sample of 74 automobiles. We know each automobile's average mileage rating and wish to test whether the overall average for the sample is 20 miles per gallon. We also assume that the population standard deviation is 6.

```
. use http://www.stata-press.com/data/r15/auto
(1978 Automobile Data)
. ztest mpg==20, sd(6)
```

One-sample z test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
mpg	74	21.2973	.6974858	6	19.93025	22.66434

```
mean = mean(mpg)                                z = 1.8600
Ho: mean = 20
Ha: mean < 20          Ha: mean != 20          Ha: mean > 20
Pr(Z < z) = 0.9686    Pr(|Z| > |z|) = 0.0629    Pr(Z > z) = 0.0314
```

The  $p$ -value for the two-sided test is 0.0629, so we do not have statistical evidence to reject the null hypothesis that the mean equals 20 at a 5% significance level, but we would reject the null hypothesis at a 10% level.

◀

## Two-sample *z* test

### ► Example 2: Two-sample *z* test using groups

We are testing the effectiveness of a new fuel additive. We run an experiment in which 12 cars are given the fuel treatment and 12 cars are not. The results of the experiment are as follows:

treated	mpg
0	20
0	23
0	21
0	25
0	18
0	17
0	18
0	24
0	20
0	24
0	23
0	19
1	24
1	25
1	21
1	22
1	23
1	18
1	17
1	28
1	24
1	27
1	21
1	23

The treated variable is coded as 1 if the car received the fuel treatment and 0 otherwise.

We can test the equality of means of the treated and untreated group by typing

```
. use http://www.stata-press.com/data/r15/fuel3
. ztest mpg, by(treated) sd(3)
```

Two-sample *z* test

Group	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
0	12	21	.8660254	3	19.30262	22.69738
1	12	22.75	.8660254	3	21.05262	24.44738
diff		-1.75	1.224745		-4.150456	.6504558

```
diff = mean(0) - mean(1)                                z = -1.4289
Ho: diff = 0
Ha: diff < 0                Ha: diff != 0                Ha: diff > 0
Pr(Z < z) = 0.0765          Pr(|Z| > |z|) = 0.1530          Pr(Z > z) = 0.9235
```

We do not have evidence to reject the null hypothesis that the means of the two groups are equal at a 5% significance level.

In the above, we assumed that the two groups have the same standard deviation of 3. If the standard deviations for the two groups are different, we can specify group-specific standard deviations in options `sd1()` and `sd2()`:

```
. ztest mpg, by(treated) sd1(2.7) sd2(3.2)
```

```
Two-sample z test
```

Group	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
0	12	21	.7794229	2.7	19.47236	22.52764
1	12	22.75	.9237604	3.2	20.93946	24.56054
diff		-1.75	1.208649		-4.118909	.6189093

```
diff = mean(0) - mean(1)
```

```
z = -1.4479
```

```
Ho: diff = 0
```

```
Ha: diff < 0
```

```
Ha: diff != 0
```

```
Ha: diff > 0
```

```
Pr(Z < z) = 0.0738
```

```
Pr(|Z| > |z|) = 0.1476
```

```
Pr(Z > z) = 0.9262
```

◀

## □ Technical note

In two-sample randomized designs, subjects will sometimes refuse the assigned treatment but still be measured for an outcome. In this case, take care to specify the group properly. You might be tempted to let *varname* contain missing where the subject refused and thus let `ztest` drop such observations from the analysis. Zelen (1979) argues that it would be better to specify that the subject belongs to the group in which he or she was randomized, even though such inclusion will dilute the measured effect.

□

## ▷ Example 3: Two-sample z test using variables

There is a second, inferior way to organize the data in the preceding example. We ran a test on 24 cars, 12 without the additive and 12 with. We now create two new variables, `mpg1` and `mpg2`.

mpg1	mpg2
20	24
23	25
21	21
25	22
18	23
17	18
18	17
24	28
20	24
24	27
23	21
19	23

This method is inferior because it suggests a connection that is not there. There is no link between the car with 20 mpg and the car with 24 mpg in the first row of the data. Each column of data could be arranged in any order. Nevertheless, if our data are organized like this, `ztest` can accommodate us.

```
. use http://www.stata-press.com/data/r15/fuel
```

```
. ztest mpg1==mpg2, unpaired sd(3)
```

```
Two-sample z test
```

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
mpg1	12	21	.8660254	3	19.30262	22.69738
mpg2	12	22.75	.8660254	3	21.05262	24.44738
diff		-1.75	1.224745		-4.150456	.6504558

```
diff = mean(mpg1) - mean(mpg2)
```

```
z = -1.4289
```

```
Ho: diff = 0
```

```
Ha: diff < 0
```

```
Ha: diff != 0
```

```
Ha: diff > 0
```

```
Pr(Z < z) = 0.0765
```

```
Pr(|Z| > |z|) = 0.1530
```

```
Pr(Z > z) = 0.9235
```

◀

## Paired z test

### ▶ Example 4

Suppose that the preceding data were actually collected by running a test on 12 cars. Each car was run once with the fuel additive and once without. Our data are stored in the same manner as in [example 3](#), but this time, there is most certainly a connection between the mpg values that appear in the same row. These come from the same car. The variables `mpg1` and `mpg2` represent mileage without and with the treatment, respectively. Suppose that the two variables have a common standard deviation of 2 and the correlation between them is 0.4.

```
. use http://www.stata-press.com/data/r15/fuel
```

```
. ztest mpg1==mpg2, sd(2) corr(0.4)
```

```
Paired z test
```

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
mpg1	12	21	.5773503	2	19.86841	22.13159
mpg2	12	22.75	.5773503	2	21.61841	23.88159
diff	12	-1.75	.6324555	2.19089	-2.98959	-.5104099

```
mean(diff) = mean(mpg1 - mpg2)
```

```
z = -2.7670
```

```
Ho: mean(diff) = 0
```

```
Ha: mean(diff) < 0
```

```
Ha: mean(diff) != 0
```

```
Ha: mean(diff) > 0
```

```
Pr(Z < z) = 0.0028
```

```
Pr(|Z| > |z|) = 0.0057
```

```
Pr(Z > z) = 0.9972
```

The *p*-value for the two-sided test is 0.0057, so we reject, for example, the null hypothesis that the two means are equal at a 5% significance level.

Equivalently, we could specify directly the standard deviation of the differences between paired observations with the `sddiff()` option:



```
. ztest mpg1==mpg2, sddiff(2.191)
```

```
Paired z test
```

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
diff	12	-1.75	.6324872	2.191	-2.989652	-.5103478

```

      mean(diff) = mean(mpg1 - mpg2)                z = -2.7669
Ho: mean(diff) = 0
Ha: mean(diff) < 0          Ha: mean(diff) != 0      Ha: mean(diff) > 0
Pr(Z < z) = 0.0028         Pr(|Z| > |z|) = 0.0057     Pr(Z > z) = 0.9972

```

◀

## Adjust for clustering

When observations are not independent and can be grouped into clusters, we need to adjust for clustering in a  $z$  test. For example, in a cluster randomized design, groups of individuals are randomized instead of individuals. To adjust for clustering, we need to specify the cluster identifier variable in the `cluster()` option. In the case of a one-sample  $z$  test, we need to also specify the intraclass correlation in the `rho()` option. In the case of a two-sample  $z$  test, we need to also specify the common population intraclass correlation in the `rho()` option or group-specific population intraclass correlations in the `rho1()` and `rho2()` options.

### ▶ Example 5: One-sample $z$ test, adjusting for clusters

Consider data on the SAT score of 75 students from 15 classes, with 5 students in each class. We want to test whether the mean verbal SAT score is different from 600. We assume a known standard deviation of 132 and a known intraclass correlation of 0.7. To perform the test, we specify the options `cluster(class)`, `rho(0.7)`, and `sd(132)`:

```
. use http://www.stata-press.com/data/r15/sat
(Fictional SAT data)
```

```
. ztest score == 600, cluster(class) rho(0.7) sd(132)
```

```

One-sample z test                Number of clusters =      15
Cluster variable: class          Cluster size           =       5
                                Intraclass corr.           =    0.7000

```

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
score	75	504.8	29.71222	132	446.5651	563.0349

```

      mean = mean(score)                z = -3.2041
Ho: mean = 600
Ha: mean < 600          Ha: mean != 600          Ha: mean > 600
Pr(Z < z) = 0.0007         Pr(|Z| > |z|) = 0.0014         Pr(Z > z) = 0.9993

```

We find statistical evidence to reject the null hypothesis of  $H_0: \mu_{\text{SAT}} = 600$  versus a two-sided alternative  $H_a: \mu_{\text{SAT}} \neq 600$  at the 5% significance level; the  $p$ -value = 0.0014 < 0.05.

◀

▷ **Example 6: Two-sample *z* test using groups, adjusting for clusters**

Consider a cluster randomized control trial that studies the effect of additional training of nurses and general practitioners in patient-centered care on the well-being and future disease risk of patients with type 2 diabetes (Kinmonth et al. [1998] and Campbell and Walters [2014]). Practices (practice) are randomly allocated to two groups—one trained to give patient-centered care (intervention group) and another trained to give routine care (comparison or control group). In our analysis, we transform the original bmi using the formula  $\ln(\text{bmi} - 14.67355)$  to obtain a variable that is approximately normally distributed, lbmi. We want to test the equality of the means of lbmi for the two groups. We assume a known common standard deviation of 0.35 and a known common intraclass correlation of 0.028.

To perform the test, we need to specify the rho(0.028) and sd(0.35) options. We also need to specify the cluster identifier practice in the cluster() option and the group identifier group in the by() option.

```
. use http://www.stata-press.com/data/r15/dcfid_trial
(BMI data from Diabetes Care from Diagnosis trial (Kinmonth et al., 1998))
.ztest lbmi, by(group) cluster(practice) rho(0.028) sd(0.35)
```

Two-sample z test  
Cluster variable: practice

<p>Group: Control</p> <p>Number of clusters =           20</p> <p>Avg. cluster size =            5.10</p> <p>CV cluster size =             0.5330</p> <p>Intraclass corr. =            0.0280</p>	<p>Group: Interv.</p> <p>Number of clusters =           18</p> <p>Avg. cluster size =            7.67</p> <p>CV cluster size =             0.5126</p> <p>Intraclass corr. =            0.0280</p>
---	---

Group	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
Control	102	2.62954	.0372502	.35	2.556531	2.702549
Interv.	138	2.749023	.0332182	.35	2.683916	2.81413
diff		-.1194831	.0499102		-.2173054	-.0216608

```
diff = mean(Control) - mean(Interv.)                    z = -2.3940
Ho: diff = 0
Ha: diff < 0                                            Ha: diff != 0                                            Ha: diff > 0
Pr(Z < z) = 0.0083                                   Pr(|Z| > |z|) = 0.0167                               Pr(Z > z) = 0.9917
```

We find statistical evidence to reject the null hypothesis of  $H_0: \mu_{\text{diff}} = 0$  versus a two-sided alternative  $H_a: \mu_{\text{diff}} \neq 0$  at the 5% significance level; the *p*-value = 0.0167 < 0.05.



**Immediate form**

▷ **Example 7: One-sample *z* test**

ztesti is like ztest, except that we specify summary statistics rather than variables as arguments. For instance, we are reading an article that reports the mean number of sunspots per month as 62.6 with a standard deviation of 15.8. We assume this standard deviation is the population standard deviation. There are 24 months of data. We wish to test whether the mean is 75:

```
. ztesti 24 62.6 15.8 75
```

```
One-sample z test
```

	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
x	24	62.6	3.225161	15.8	56.2788	68.9212

```

mean = mean(x)                                z = -3.8448
Ho: mean = 75
Ha: mean < 75                                Ha: mean != 75                                Ha: mean > 75
Pr(Z < z) = 0.0001                            Pr(|Z| > |z|) = 0.0001                            Pr(Z > z) = 0.9999

```

◀

## ▶ Example 8: Two-sample $z$ test

There is no immediate form of `ztest` with paired data because the test is also a function of the covariance, a number unlikely to be reported in any published source. For unpaired data, however, we might type

```
. ztesti 20 20 5 32 15 4
```

```
Two-sample z test
```

	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
x	20	20	1.118034	5	17.80869	22.19131
y	32	15	.7071068	4	13.6141	16.3859
diff		5	1.322876		2.407211	7.592789

```

diff = mean(x) - mean(y)                        z = 3.7796
Ho: diff = 0
Ha: diff < 0                                Ha: diff != 0                                Ha: diff > 0
Pr(Z < z) = 0.9999                            Pr(|Z| > |z|) = 0.0002                            Pr(Z > z) = 0.0001

```

◀

## Stored results

One-sample `ztest` and `ztesti` store the following in `r()`:

Scalars

<code>r(N)</code>	sample size
<code>r(mu)</code>	sample mean
<code>r(sd)</code>	standard deviation
<code>r(se)</code>	standard error
<code>r(lb)</code>	lower confidence bound of one-sample mean
<code>r(ub)</code>	upper confidence bound of one-sample mean
<code>r(z)</code>	$z$ statistic
<code>r(p_l)</code>	lower one-sided $p$ -value
<code>r(p)</code>	two-sided $p$ -value
<code>r(p_u)</code>	upper one-sided $p$ -value
<code>r(level)</code>	confidence level

Cluster-adjusted one-sample `ztest` also stores the following in `r()`:

Scalars

<code>r(K)</code>	number of clusters $K$
<code>r(M)</code>	cluster size $M$
<code>r(rho)</code>	intraclass correlation
<code>r(CV_cluster)</code>	coefficient of variation for cluster sizes

Two-sample `ztest` and `ztesti` store the following in `r()`:

Scalars

<code>r(N1)</code>	sample size of population one
<code>r(N2)</code>	sample size of population two
<code>r(mu1)</code>	sample mean for population one
<code>r(mu2)</code>	sample mean for population two
<code>r(mu_diff)</code>	difference of means
<code>r(corr)</code>	correlation between paired observations; if the <code>corr()</code> option is specified
<code>r(sd)</code>	common standard deviation
<code>r(sd1)</code>	standard deviation for population one
<code>r(sd2)</code>	standard deviation for population two
<code>r(sd_diff)</code>	standard deviation of the differences between paired observations
<code>r(se1)</code>	standard error of population-one sample mean
<code>r(se2)</code>	standard error of population-two sample mean
<code>r(se_diff)</code>	standard error of the difference of means
<code>r(lb1)</code>	lower confidence bound of population-one sample mean
<code>r(ub1)</code>	upper confidence bound of population-one sample mean
<code>r(lb2)</code>	lower confidence bound of population-two sample mean
<code>r(ub2)</code>	upper confidence bound of population-two sample mean
<code>r(lb_diff)</code>	lower confidence bound of the difference of means
<code>r(ub_diff)</code>	upper confidence bound of the difference of means
<code>r(z)</code>	$z$ statistic
<code>r(p_l)</code>	lower one-sided $p$ -value
<code>r(p)</code>	two-sided $p$ -value
<code>r(p_u)</code>	upper one-sided $p$ -value
<code>r(level)</code>	confidence level

Cluster-adjusted two-sample `ztest` using the `by()` option also stores the following in `r()`:

Scalars

<code>r(K1)</code>	population-one number of clusters $K_1$
<code>r(K2)</code>	population-two number of clusters $K_2$
<code>r(M1)</code>	population-one cluster size $M_1$
<code>r(M2)</code>	population-two cluster size $M_2$
<code>r(rho)</code>	common intraclass correlation
<code>r(rho1)</code>	population-one intraclass correlation
<code>r(rho2)</code>	population-two intraclass correlation
<code>r(CV_cluster1)</code>	population-one coefficient of variation for cluster sizes
<code>r(CV_cluster2)</code>	population-two coefficient of variation for cluster sizes

## Methods and formulas

Methods and formulas are presented under the following headings:

*One-sample  $z$  test*

*Two-sample unpaired  $z$  test*

*Paired  $z$  test*

For all the tests below, the test statistic  $z$  is distributed as standard normal, and the  $p$ -value is computed as

$$p = \begin{cases} 1 - \Phi(z) & \text{for an upper one-sided test} \\ \Phi(z) & \text{for a lower one-sided test} \\ 2\{1 - \Phi(|z|)\} & \text{for a two-sided test} \end{cases}$$

where  $\Phi(\cdot)$  is the cdf of a standard normal distribution and  $|z|$  is an absolute value of  $z$ .

Also see, for instance, [Hoel \(1984, 140–161\)](#), [Dixon and Massey \(1983, 100–130\)](#), and [Tamhane and Dunlop \(2000, 237–290\)](#) for more information about  $z$  tests.

## One-sample $z$ test

Suppose that we observe a random sample  $x_1, x_2, \dots, x_n$  of size  $n$ , which follows a normal distribution with mean  $\mu$  and standard deviation  $\sigma$ . We are interested in testing the null hypothesis  $H_0: \mu = \mu_0$  versus the two-sided alternative hypothesis  $H_a: \mu \neq \mu_0$ , the upper one-sided alternative  $H_a: \mu > \mu_0$ , or the lower one-sided alternative  $H_a: \mu < \mu_0$ . Assuming a known standard deviation  $\sigma$ , we use the following test statistic,

$$z = \frac{(\bar{x} - \mu_0)}{s}$$

where  $\bar{x} = (\sum_{i=1}^n x_i)/n$  is the sample mean and  $s = \sigma/\sqrt{n}$  is the standard error of  $\bar{x}$ .

The  $100(1 - \alpha)\%$  confidence interval for  $\bar{x}$  is given by

$$\bar{x} \pm z_{1-\alpha/2}s$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ th quantile of the standard normal distribution.

With clustered data, suppose that there are  $K$  clusters. The  $i$ th cluster of size  $M_i$  contains the observations  $x_{i1}, x_{i2}, \dots, x_{iM_i}$ , such that  $n = \sum_{i=1}^K M_i$  and  $\bar{x} = \frac{1}{n} \sum_{i=1}^K \sum_{j=1}^{M_i} x_{ij}$ . Let  $\rho$  be the intraclass correlation. Following [Ahn, Heo, and Zhang \(2015\)](#), we assume that the cluster sizes  $M_i$  are independent and identically distributed. Let  $C_{\text{adj}}$  be the adjustment to the standard error for clustered data,

$$C_{\text{adj}} = \sqrt{\sum_{i=1}^K M_i \{1 + \rho(M_i - 1)\} / n}$$

such that  $s_{\text{cl}} = C_{\text{adj}}s$ .

$C_{\text{adj}}$  can be equivalently written as

$$C_{\text{adj}} = \sqrt{1 + \rho(\bar{M} - 1) + \rho\bar{M}\text{CV}_{\text{cl}}^2}$$

where  $\bar{M} = \sum_{i=1}^K M_i/K$  is the average cluster size and  $\text{CV}_{\text{cl}}$  is the coefficient of variation for cluster sizes:

$$\text{CV}_{\text{cl}} = \frac{\sqrt{\sum_{i=1}^K (M_i - \bar{M})^2 / K}}{\bar{M}}$$

To adjust the test statistic  $z$  and the confidence interval for clustering, replace  $s$  with  $s_{\text{cl}}$  in the corresponding formulas. In the presence of clustering, the test statistic  $z$  is asymptotically normally distributed conditional on the empirical distribution of  $M_i$ 's.

## Two-sample unpaired $z$ test

Suppose that we observe a random sample  $x_{11}, x_{12}, \dots, x_{1n_1}$  of size  $n_1$ , which follows a normal distribution with mean  $\mu_1$  and standard deviation  $\sigma_1$ , and another random sample  $x_{21}, x_{22}, \dots, x_{2n_2}$  of size  $n_2$ , which follows a normal distribution with mean  $\mu_2$  and standard deviation  $\sigma_2$ . We are interested in testing the null hypothesis  $H_0: \mu_2 = \mu_1$  versus the two-sided alternative hypothesis  $H_a: \mu_2 \neq \mu_1$ , the upper one-sided alternative  $H_a: \mu_2 > \mu_1$ , or the lower one-sided alternative  $H_a: \mu_2 < \mu_1$ . Assuming known standard deviations  $\sigma_1$  and  $\sigma_2$ , we use the following test statistic,

$$z = \frac{\bar{x}_2 - \bar{x}_1}{\sqrt{s_1^2 + s_2^2}}$$

where  $\bar{x}_1 = (\sum_{i=1}^{n_1} x_{1i})/n_1$  and  $\bar{x}_2 = (\sum_{i=1}^{n_2} x_{2i})/n_2$  are the two sample means and  $s_1 = \sigma_1/\sqrt{n_1}$  and  $s_2 = \sigma_2/\sqrt{n_2}$  are the corresponding two standard errors.

The  $100(1 - \alpha)\%$  confidence intervals for  $\bar{x}_1$  and  $\bar{x}_2$  are given by

$$\begin{aligned} \bar{x}_1 \pm z_{1-\alpha/2} s_1 \\ \bar{x}_2 \pm z_{1-\alpha/2} s_2 \end{aligned}$$

where  $z_{1-\alpha/2}$  is the  $(1 - \alpha/2)$ th quantile of the standard normal distribution.

The  $100(1 - \alpha)\%$  confidence interval for  $\bar{x}_1 - \bar{x}_2$  is given by

$$\bar{x}_1 - \bar{x}_2 \pm z_{1-\alpha/2} \sqrt{s_1^2 + s_2^2}$$

With clustered data, similar to the discussion for the one-sample test, suppose that population one has  $K_1$  clusters and population two has  $K_2$  clusters. Let  $\rho_1$  and  $\rho_2$  be the intraclass correlations,  $\bar{M}_1$  and  $\bar{M}_2$  be the average cluster sizes,  $\bar{x}_1 = (1/n_1) \sum_{i=1}^{K_1} \sum_{j=1}^{M_{1i}} x_{1ij}$  and  $\bar{x}_2 = (1/n_2) \sum_{i=1}^{K_2} \sum_{j=1}^{M_{2i}} x_{2ij}$  be the sample means, and  $\text{CV}_{\text{cl},1}$  and  $\text{CV}_{\text{cl},2}$  be the coefficients of variation for cluster sizes for population one and population two. Let  $s_{1,\text{cl}} = C_{\text{adj},1} s_1$  and  $s_{2,\text{cl}} = C_{\text{adj},2} s_2$  be the standard errors of the population-specific sample means adjusted for clustered data, where the population-specific adjustment factors are defined as described for the one-sample test. To adjust the two-sample test statistic and the confidence intervals for clustering, replace  $s_1$  with  $s_{1,\text{cl}}$  and  $s_2$  with  $s_{2,\text{cl}}$  in the corresponding formulas.

## Paired $z$ test

Some experiments have paired observations (also known as matched observations, correlated pairs, or permanent components). Consider a sequence of  $n$  paired observations denoted by  $x_{ij}$  for subjects  $i = 1, 2, \dots, n$  and groups  $j = 1, 2$ . An individual observation corresponds to the pair  $(x_{i1}, x_{i2})$ , and inference is made on the differences within the pairs. Let  $\mu_d = \mu_2 - \mu_1$  denote the mean difference, where  $\mu_j$  is the population mean of group  $j$ , and let  $D_i = x_{i2} - x_{i1}$  denote the difference between individual observations.  $D_i$  follows a normal distribution with mean  $\mu_2 - \mu_1$  and standard deviation  $\sigma_d$ , where  $\sigma_d = \sqrt{\sigma_1^2 + \sigma_2^2 - 2\rho_{\text{pair}}\sigma_1\sigma_2}$ ,  $\sigma_j$  is the population standard deviation of group  $j$  and  $\rho_{\text{pair}}$  is the correlation between paired observations.

We are interested in testing the null hypothesis  $H_0: \mu_2 = \mu_1$  versus the two-sided alternative hypothesis  $H_a: \mu_2 \neq \mu_1$ , the upper one-sided alternative  $H_a: \mu_2 > \mu_1$ , or the lower one-sided alternative  $H_a: \mu_2 < \mu_1$ . Assuming the standard deviation of the differences  $\sigma_d$  is known, we use the following test statistic,

$$z = \frac{\bar{d}}{s_d}$$

where  $\bar{d} = (\sum_{i=1}^n D_i)/n$  is the sample mean of the differences between paired observations and  $s_d = \sigma_d/\sqrt{n}$  is the standard error of  $\bar{d}$ .

The  $100(1 - \alpha)\%$  confidence interval for  $\bar{d}$  is given by

$$\bar{d} \pm z_{1-\alpha/2} s_d$$

## References

- Ahn, C., M. Heo, and S. Zhang. 2015. *Sample Size Calculations for Clustered and Longitudinal Outcomes in Clinical Research*. Boca Raton, FL: CRC Press.
- Campbell, M. J., and S. J. Walters. 2014. *How to Design, Analyse and Report Cluster Randomised Trials in Medicine and Health Related Research*. Chichester, UK: Wiley.
- Dixon, W. J., and F. J. Massey, Jr. 1983. *Introduction to Statistical Analysis*. 4th ed. New York: McGraw-Hill.
- Hoel, P. G. 1984. *Introduction to Mathematical Statistics*. 5th ed. New York: Wiley.
- Kinmonth, A. L., A. Woodcock, S. Griffin, N. Spiegel, and M. J. Campbell. 1998. Randomised controlled trial of patient centred care of diabetes in general practice: Impact on current wellbeing and future disease risk. *British Medical Journal* 317: 1202–1208.
- Tamhane, A. C., and D. D. Dunlop. 2000. *Statistics and Data Analysis: From Elementary to Intermediate*. Upper Saddle River, NJ: Prentice Hall.
- Zelen, M. 1979. A new design for randomized clinical trials. *New England Journal of Medicine* 300: 1242–1245.

## Also see

- [R] **ci** — Confidence intervals for means, proportions, and variances
- [R] **esize** — Effect size based on mean comparison
- [R] **mean** — Estimate means
- [R] **oneway** — One-way analysis of variance
- [R] **ttest** —  $t$  tests (mean-comparison tests)
- [MV] **hotelling** — Hotelling's T-squared generalized means test
- [PSS] **power onemean** — Power analysis for a one-sample mean test
- [PSS] **power onemean, cluster** — Power analysis for a one-sample mean test, CRD
- [PSS] **power pairedmeans** — Power analysis for a two-sample paired-means test
- [PSS] **power twomeans** — Power analysis for a two-sample means test
- [PSS] **power twomeans, cluster** — Power analysis for a two-sample means test, CRD