

example 20 — Two-factor measurement model by group
[Description](#)[Remarks and examples](#)[Reference](#)[Also see](#)

Description

Below we demonstrate `sem`'s `group()` option, which allows fitting models in which path coefficients and covariances differ across groups of the data, such as for males and females. We use the following data:

```
. use http://www.stata-press.com/data/r15/sem_2fmmby
(two-factor CFA)
. ssd describe

Summary statistics data from
http://www.stata-press.com/data/r15/sem_2fmmby.dta
  obs:          385                two-factor CFA
  vars:         16                25 May 2016 11:11
                                   (_dta has notes)
```

variable name	variable label
phyab1	Physical ability 1
phyab2	Physical ability 2
phyab3	Physical ability 3
phyab4	Physical ability 4
appear1	Appearance 1
appear2	Appearance 2
appear3	Appearance 3
appear4	Appearance 4
peerrel1	Relationship w/ peers 1
peerrel2	Relationship w/ peers 2
peerrel3	Relationship w/ peers 3
peerrel4	Relationship w/ peers 4
parrel1	Relationship w/ parent 1
parrel2	Relationship w/ parent 2
parrel3	Relationship w/ parent 3
parrel4	Relationship w/ parent 4

```
Group variable: grade (2 groups)
```

```
Obs. by group: 134, 251
```

```
. notes
```

```
_dta:
```

1. Summary statistics data from Marsh, H. W. and Hocevar, D., 1985, "Application of confirmatory factor analysis to the study of self-concept: First- and higher order factor models and their invariance across groups", *_Psychological Bulletin_*, 97: 562-582.
2. Summary statistics based on 134 students in grade 4 and 251 students in grade 5 from Sydney, Australia.
3. Group 1 is grade 4, group 2 is grade 5.
4. Data collected using the Self-Description Questionnaire and includes sixteen subscales designed to measure nonacademic status: four intended to measure physical ability, four intended to measure physical appearance, four intended to measure relations with peers, and four intended to measure relations with parents.

Remarks and examples

Remarks are presented under the following headings:

Background

Fitting the model with all the data


Fitting the model with the `group()` option

Fitting the model with the Builder

Background

See [SEM] intro 6 for background on `sem`'s `group()` option.

We will fit the model



which in command syntax can be written as

```
(Peer -> peerrel1 peerrel2 peerrel3 peerrel4)   ///
(Par  -> parrel1  parrel2  parrel3  parrel4)
```

We are using the same data used in [SEM] example 15, but we are using more of the data and fitting a different model. To remind you, those data were collected from students in grade 5. The dataset we are using, however, has data for students from grade 4 and from grade 5, which was created in [SEM] example 19. We have the following observed variables:

1. Four measures of physical ability.
2. Four measures of appearance.
3. Four measures of quality of relationship with peers.
4. Four measures of quality of relationship with parents.

In this example, we will consider solely the measurement problem, and include only the measurement variables for the two kinds of relationship quality. We are going to treat quality of relationship with peers as measures of underlying factor `Peer` and quality of relationship with parents as measures of underlying factor `Par`.

Below we will

1. Fit the model with all the data. This amounts to assuming that the students in grades 4 and 5 are identical in terms of this measurement problem.
2. Fit the model with `sem`'s `group()` option, which will constrain some parameters to be the same for students in grades 4 and 5 and leave free of constraint the others.

Fitting the model with all the data

Throughout this example, we want you to appreciate that we are using SSD and that matters not at all. Not one command would have a different syntax or option, or produce a different result, if we had the real data.

We begin by fitting the model with all the data:

```
. sem (Peer -> peerrel1 peerrel2 peerrel3 peerrel4)
>      (Par -> parrel1 parrel2 parrel3 parrel4)
Endogenous variables
Measurement: peerrel1 peerrel2 peerrel3 peerrel4 parrel1 parrel2 parrel3 parrel4
Exogenous variables
Latent:      Peer Par
Fitting target model:
Iteration 0:  log likelihood = -5559.545
Iteration 1:  log likelihood = -5558.609
Iteration 2:  log likelihood = -5558.6017
Iteration 3:  log likelihood = -5558.6017
```

4 example 20 — Two-factor measurement model by group

```

Structural equation model          Number of obs   =       385
Estimation method = ml
Log likelihood = -5558.6017
( 1) [peerrel1]Peer = 1
( 2) [parrel1]Par = 1

```

	OIM				[95% Conf. Interval]	
	Coef.	Std. Err.	z	P> z		
Measurement						
peerrel1						
Peer	1 (constrained)					
_cons	8.681221	.0937197	92.63	0.000	8.497534	8.864908
peerrel2						
Peer	1.113865	.09796	11.37	0.000	.9218666	1.305863
_cons	7.828623	.1037547	75.45	0.000	7.625268	8.031979
peerrel3						
Peer	1.42191	.114341	12.44	0.000	1.197806	1.646014
_cons	7.359896	.1149905	64.00	0.000	7.134519	7.585273
peerrel4						
Peer	1.204146	.0983865	12.24	0.000	1.011312	1.39698
_cons	8.150779	.1023467	79.64	0.000	7.950183	8.351375
parrel1						
Par	1 (constrained)					
_cons	9.339558	.0648742	143.96	0.000	9.212407	9.46671
parrel2						
Par	1.112383	.1378687	8.07	0.000	.8421655	1.382601
_cons	9.220494	.0742356	124.21	0.000	9.074994	9.365993
parrel3						
Par	2.037924	.204617	9.96	0.000	1.636882	2.438966
_cons	8.676961	.088927	97.57	0.000	8.502667	8.851255
parrel4						
Par	1.52253	.1536868	9.91	0.000	1.221309	1.82375
_cons	9.045247	.0722358	125.22	0.000	8.903667	9.186826
var(e.peer~1)	1.809309	.1596546			1.521956	2.150916
var(e.peer~2)	2.193804	.194494			1.843884	2.610129
var(e.peer~3)	1.911874	.214104			1.535099	2.381126
var(e.peer~4)	1.753037	.1749613			1.441575	2.131792
var(e.parr~1)	1.120333	.0899209			.9572541	1.311193
var(e.parr~2)	1.503003	.1200739			1.285162	1.757769
var(e.parr~3)	.9680081	.1419777			.7261617	1.290401
var(e.parr~4)	.8498834	.0933687			.685245	1.054078
var(Peer)	1.572294	.2255704			1.186904	2.082822
var(Par)	.5000022	.093189			.3469983	.7204709
cov(Peer,Par)	.4226706	.0725253	5.83	0.000	.2805236	.5648176

LR test of model vs. saturated: chi2(19) = 28.19, Prob > chi2 = 0.0798

Note:

1. We are using SSD with data for two separate groups. There is no hint of that in the output above because sem combined the summary statistics and produced overall results just as if we had the real data.

Fitting the model with the group() option

```

. sem (Peer -> peerrel1 peerrel2 peerrel3 peerrel4)
>     (Par -> parrel1 parrel2 parrel3 parrel4), group(grade)

Endogenous variables
Measurement: peerrel1 peerrel2 peerrel3 peerrel4 parrel1 parrel2 parrel3 parrel4
Exogenous variables
Latent:      Peer Par
Fitting target model:
Iteration 0: log likelihood = -13049.77 (not concave)
Iteration 1: log likelihood = -10819.682 (not concave)
Iteration 2: log likelihood = -8873.4568 (not concave)
Iteration 3: log likelihood = -6119.7114 (not concave)
Iteration 4: log likelihood = -5949.354 (not concave)
Iteration 5: log likelihood = -5775.6085 (not concave)
Iteration 6: log likelihood = -5713.9178 (not concave)
Iteration 7: log likelihood = -5638.1208 (not concave)
Iteration 8: log likelihood = -5616.6335 (not concave)
Iteration 9: log likelihood = -5595.7507 (not concave)
Iteration 10: log likelihood = -5589.9802 (not concave)
Iteration 11: log likelihood = -5578.8701 (not concave)
Iteration 12: log likelihood = -5574.0162 (not concave)
Iteration 13: log likelihood = -5568.0786
Iteration 14: log likelihood = -5551.7349
Iteration 15: log likelihood = -5544.0052
Iteration 16: log likelihood = -5542.7113
Iteration 17: log likelihood = -5542.6775
Iteration 18: log likelihood = -5542.6774

Structural equation model
Grouping variable = grade
Estimation method = ml
Log likelihood = -5542.6774

Number of obs = 385
Number of groups = 2

( 1) [peerrel1]1bn.grade#c.Peer = 1
( 2) [peerrel2]1bn.grade#c.Peer - [peerrel2]2.grade#c.Peer = 0
( 3) [peerrel3]1bn.grade#c.Peer - [peerrel3]2.grade#c.Peer = 0
( 4) [peerrel4]1bn.grade#c.Peer - [peerrel4]2.grade#c.Peer = 0
( 5) [parrel1]1bn.grade#c.Par = 1
( 6) [parrel2]1bn.grade#c.Par - [parrel2]2.grade#c.Par = 0
( 7) [parrel3]1bn.grade#c.Par - [parrel3]2.grade#c.Par = 0
( 8) [parrel4]1bn.grade#c.Par - [parrel4]2.grade#c.Par = 0
( 9) [peerrel1]1bn.grade - [peerrel1]2.grade = 0
(10) [peerrel2]1bn.grade - [peerrel2]2.grade = 0
(11) [peerrel3]1bn.grade - [peerrel3]2.grade = 0
(12) [peerrel4]1bn.grade - [peerrel4]2.grade = 0
(13) [parrel1]1bn.grade - [parrel1]2.grade = 0
(14) [parrel2]1bn.grade - [parrel2]2.grade = 0
(15) [parrel3]1bn.grade - [parrel3]2.grade = 0
(16) [parrel4]1bn.grade - [parrel4]2.grade = 0
(17) [peerrel1]2.grade#c.Peer = 1
(18) [parrel1]2.grade#c.Par = 1
(19) [/]mean(Peer)#1bn.grade = 0
(20) [/]mean(Par)#1bn.grade = 0

```

6 example 20 — Two-factor measurement model by group

Group	: 1	Number of obs				=	134
	OIM		z	P> z	[95% Conf. Interval]		
	Coef.	Std. Err.					
Measurement							
peerrel1							
Peer	1 (constrained)						
_cons	8.466539	.1473448	57.46	0.000	8.177748	8.755329	
peerrel2							
Peer	1.109234	.0975279	11.37	0.000	.9180833	1.300385	
_cons	7.589872	.1632145	46.50	0.000	7.269977	7.909766	
peerrel3							
Peer	1.409361	.1138314	12.38	0.000	1.186256	1.632467	
_cons	7.056996	.1964299	35.93	0.000	6.672001	7.441992	
peerrel4							
Peer	1.195982	.0980272	12.20	0.000	1.003852	1.388112	
_cons	7.89358	.169158	46.66	0.000	7.562036	8.225123	
parrel1							
Par	1 (constrained)						
_cons	9.368654	.0819489	114.32	0.000	9.208037	9.529271	
parrel2							
Par	1.104355	.1369365	8.06	0.000	.8359649	1.372746	
_cons	9.287629	.0903296	102.82	0.000	9.110587	9.464672	
parrel3							
Par	2.05859	.2060583	9.99	0.000	1.654723	2.462457	
_cons	8.741898	.136612	63.99	0.000	8.474144	9.009653	
parrel4							
Par	1.526706	.1552486	9.83	0.000	1.222424	1.830987	
_cons	9.096609	.1061607	85.69	0.000	8.888538	9.30468	
mean(Peer)	0 (constrained)						
mean(Par)	0 (constrained)						
var(e.peer~1)	1.824193	.2739446			1.359074	2.448489	
var(e.peer~2)	2.236974	.3310875			1.673699	2.989817	
var(e.peer~3)	1.907009	.3383293			1.346908	2.700023	
var(e.peer~4)	1.639881	.272764			1.18367	2.271925	
var(e.parr~1)	.9669121	.1302489			.7425488	1.259067	
var(e.parr~2)	.9683878	.133192			.7395628	1.268012	
var(e.parr~3)	.8377567	.1986089			.526407	1.333258	
var(e.parr~4)	.8343032	.1384649			.6026352	1.15503	
var(Peer)	2.039297	.3784544			1.41747	2.933912	
var(Par)	.4492996	.1011565			.2889976	.6985183	
cov(Peer,Par)	.5012091	.1193333	4.20	0.000	.2673201	.7350982	

Group	: 2		Number of obs		=	251	
	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]		
Measurement							
peerrel1							
Peer	1	(constrained)					
_cons	8.466539	.1473448	57.46	0.000	8.177748	8.755329	
peerrel2							
Peer	1.109234	.0975279	11.37	0.000	.9180833	1.300385	
_cons	7.589872	.1632145	46.50	0.000	7.269977	7.909766	
peerrel3							
Peer	1.409361	.1138314	12.38	0.000	1.186256	1.632467	
_cons	7.056996	.1964299	35.93	0.000	6.672001	7.441992	
peerrel4							
Peer	1.195982	.0980272	12.20	0.000	1.003852	1.388112	
_cons	7.89358	.169158	46.66	0.000	7.562036	8.225123	
parrel1							
Par	1	(constrained)					
_cons	9.368654	.0819489	114.32	0.000	9.208037	9.529271	
parrel2							
Par	1.104355	.1369365	8.06	0.000	.8359649	1.372746	
_cons	9.287629	.0903296	102.82	0.000	9.110587	9.464672	
parrel3							
Par	2.05859	.2060583	9.99	0.000	1.654723	2.462457	
_cons	8.741898	.136612	63.99	0.000	8.474144	9.009653	
parrel4							
Par	1.526706	.1552486	9.83	0.000	1.222424	1.830987	
_cons	9.096609	.1061607	85.69	0.000	8.888538	9.30468	
mean(Peer)	.3296841	.1570203	2.10	0.036	.02193	.6374382	
mean(Par)	-.0512439	.0818255	-0.63	0.531	-.211619	.1091313	
var(e.peer~1)	1.773813	.1889104			1.439644	2.185549	
var(e.peer~2)	2.165228	.2321565			1.75484	2.671589	
var(e.peer~3)	1.950679	.2586196			1.504298	2.529516	
var(e.peer~4)	1.822448	.2151827			1.445942	2.296992	
var(e.parr~1)	1.213159	.1192634			1.000547	1.470949	
var(e.parr~2)	1.79031	.1747374			1.478596	2.167739	
var(e.parr~3)	1.015707	.1713759			.7297073	1.4138	
var(e.parr~4)	.8599648	.1165865			.6592987	1.121706	
var(Peer)	1.307976	.2061581			.9603661	1.781406	
var(Par)	.5201696	.1029353			.3529413	.7666329	
cov(Peer,Par)	.3867156	.079455	4.87	0.000	.2309867	.5424445	

LR test of model vs. saturated: $\chi^2(50) = 61.91$, Prob > $\chi^2 = 0.1204$

Notes:

1. In *Which parameters vary by default, and which do not* in [SEM] intro 6, we wrote that, generally speaking, when we specify `group(groupvar)`, the measurement part of the model is constrained by default to be the same across the groups, whereas the remaining parts will have separate parameters for each group.

More precisely, we revealed that `sem` classifies each parameter into one of nine classes, which are the following:

Class description	Class name
1. structural coefficients	<code>scoef</code>
2. structural intercepts	<code>scons</code>
3. measurement coefficients	<code>mcoef</code>
4. measurement intercepts	<code>mcons</code>
5. covariances of structural errors	<code>serrvar</code>
6. covariances of measurement errors	<code>merrvar</code>
7. covariances between structural and measurement errors	<code>smerrcov</code>
8. means of exogenous variables	<code>meanex</code> (*)
9. covariances of exogenous variables	<code>covex</code> (*)
10. all the above	<code>all</code> (*)
11. none of the above	<code>none</code>

(*) Be aware that classes 8, 9, and 10 (`meanex`, `covex`, and `all`) exclude the observed exogenous variables—include only the latent exogenous variables—unless you specify option `noxconditional` or the `noxconditional` option is otherwise implied; see [SEM] [sem option noxconditional](#). This is what you would desire in most cases.

By default, classes 3 and 4 are constrained to be equal and the rest are allowed to vary.

2. Thus you might expect that most of the parameters of our model would have been left unconstrained until you remember that we are fitting a measurement model. That is why `sem` listed 20 constraints at the top of the estimation results. Some of the constraints are substantive and some are normalization.

3. In the output, we have a separate table of parameter estimates for each level of `grade`.

In our data, group 1 corresponds with students in grade 4, and group 2 corresponds with students in grade 5.

4. It may surprise you that the output contains estimates for the means of the latent variables. Usually, `sem` does not report this.

Usually, you are running on only one group of data and those means cannot be estimated, at least not without additional identifying constraints. When you are running on two or more groups, the means for all the groups except one can be estimated.

In [SEM] [example 21](#), we use `estat ggoof` to evaluate goodness of fit group by group.

In [SEM] [example 22](#), we use `estat ginvariant` to test whether parameters that are constrained across groups should not be and whether parameters that are not constrained could be.

In [SEM] [example 23](#), we show how to constrain the parameters we choose to be equal across groups.

Fitting the model with the Builder

Use the diagram above for reference.

1. Open the dataset.


In the Command window, type

```
. use http://www.stata-press.com/data/r15/sem_2fmmbv
```

2. Open a new Builder diagram.

Select menu item **Statistics > SEM (structural equation modeling) > Model building and estimation**.

3. Create the measurement component for relationships with peers.


Select the Add measurement component tool, , and then click in the diagram about halfway down from the top and about one-third of the way in from the left.

In the resulting dialog box,

- a. change the *Latent variable name* to **Peer**;
- b. select **peerrel1**, **peerrel2**, **peerrel3**, and **peerrel4** by using the *Measurement variables control*;
- c. select **Left** in the *Measurement direction control*;
- d. click on **OK**.

If you wish, move the component by clicking on any variable and dragging it.

4. Create the measurement component for relationships with parents.


Select the Add measurement component tool, , and then click in the diagram about halfway down from the top and about one-third of the way in from the right.

In the resulting dialog box,


- a. change the *Latent variable name* to **Par**;
- b. select **parrel1**, **parrel2**, **parrel3**, and **parrel4** by using the *Measurement variables control*;
- c. select **Right** in the *Measurement direction control*;
- d. click on **OK**.

If you wish, move the component by clicking on any variable and dragging it.


5. Correlate the latent variables.

- a. Select the Add covariance tool, .
- b. Click in the upper-right quadrant of the **Peer** oval (it will highlight when you hover over it), and drag a covariance to the upper-left quadrant of the **Par** oval (it will highlight when you can release to connect the covariance).

6. Clean up.

If you do not like where a covariance has been connected to its variable, use the Select tool, , to click on the covariance, and then simply click on where it connects to an oval and drag the endpoint. You can also change the bow of the covariance by dragging the control point that extends from one end of the selected covariance.

7. Estimate.

Click on the **Estimate** button, , in the Standard Toolbar.

In the resulting dialog box, do the following:

- a. Select the **Group** tab.
- b. Select the *Group analysis* radio button. The variable `grade` should appear in the *Group variable* control.
- c. Click on **OK**.
- d. In the Standard Toolbar, use the *Group* control to toggle between results for group 1 and group 2.

You can open a completed diagram in the Builder by typing

```
. webgetsem sem_2fmmby
```

Reference

Acock, A. C. 2013. *Discovering Structural Equation Modeling Using Stata*. Rev. ed. College Station, TX: Stata Press.

Also see

[SEM] **example 3** — Two-factor measurement model

[SEM] **example 19** — Creating multiple-group summary statistics data

[SEM] **example 21** — Group-level goodness of fit

[SEM] **example 22** — Testing parameter equality across groups

[SEM] **example 23** — Specifying parameter constraints across groups

[SEM] **intro 6** — Comparing groups

[SEM] **sem** — Structural equation model estimation command

[SEM] **sem group options** — Fitting models on different groups