

ameans — Arithmetic, geometric, and harmonic means

Syntax	Menu	Description	Options
Remarks and examples	Stored results	Methods and formulas	Acknowledgments
References	Also see		

Syntax

```
ameans [varlist] [if] [in] [weight] [, options]
```

<i>options</i>	Description
----------------	-------------

Main

<code>add(#)</code>	add # to each variable in <i>varlist</i>
<code>only</code>	add # only to variables with nonpositive values
<code>level(#)</code>	set confidence level; default is <code>level(95)</code>

`by` is allowed; see [\[D\] by](#).

`aweight`s and `fweight`s are allowed; see [\[U\] 11.1.6 weight](#).

Menu

Statistics > Summaries, tables, and tests > Summary and descriptive statistics > Arith./geometric/harmonic means

Description

`ameans` computes the arithmetic, geometric, and harmonic means, with their corresponding confidence intervals, for each variable in *varlist* or for all the variables in the data if *varlist* is not specified. `gmeans` and `hmeans` are synonyms for `ameans`.

If you simply want arithmetic means and corresponding confidence intervals, see [\[R\] ci](#).

Options

Main

`add(#)` adds the value # to each variable in *varlist* before computing the means and confidence intervals. This option is useful when analyzing variables with nonpositive values.

`only` modifies the action of the `add(#)` option so that it adds # only to variables with at least one nonpositive value.

`level(#)` specifies the confidence level, as a percentage, for confidence intervals. The default is `level(95)` or as set by `set level`; see [\[U\] 20.7 Specifying the width of confidence intervals](#).

Remarks and examples

▷ Example 1

We have a dataset containing 8 observations on a variable named `x`. The eight values are 5, 4, -4, -5, 0, 0, *missing*, and 7.

```
. ameans x
```

Variable	Type	Obs	Mean	[95% Conf. Interval]	
x	Arithmetic	7	1	-3.204405	5.204405
	Geometric	3	5.192494	2.57899	10.45448
	Harmonic	3	5.060241	3.023008	15.5179

```
. ameans x, add(5)
```

Variable	Type	Obs	Mean	[95% Conf. Interval]	
x	Arithmetic	7	6	1.795595	10.2044 *
	Geometric	6	5.477226	2.1096	14.22071 *
	Harmonic	6	3.540984	.	. *

(*) 5 was added to the variables prior to calculating the results. Missing values in confidence intervals for harmonic mean indicate that confidence interval is undefined for corresponding variables. Consult Reference Manual for details.

The number of observations displayed for the arithmetic mean is the number of nonmissing observations. The number of observations displayed for the geometric and harmonic means is the number of nonmissing, positive observations. Specifying the `add(5)` option produces 3 more positive observations. The confidence interval for the harmonic mean is not reported; see [Methods and formulas](#) below. ◀

Video example

[Descriptive statistics in Stata](#)

Stored results

`ameans` stores the following in `r()`:

Scalars

<code>r(N)</code>	number of nonmissing observations; used for arithmetic mean
<code>r(N_pos)</code>	number of nonmissing positive observations; used for geometric and harmonic means
<code>r(mean)</code>	arithmetic mean
<code>r(lb)</code>	lower bound of confidence interval for arithmetic mean
<code>r(ub)</code>	upper bound of confidence interval for arithmetic mean
<code>r(Var)</code>	variance of untransformed data
<code>r(mean_g)</code>	geometric mean
<code>r(lb_g)</code>	lower bound of confidence interval for geometric mean
<code>r(ub_g)</code>	upper bound of confidence interval for geometric mean
<code>r(Var_g)</code>	variance of $\ln x_i$
<code>r(mean_h)</code>	harmonic mean
<code>r(lb_h)</code>	lower bound of confidence interval for harmonic mean
<code>r(ub_h)</code>	upper bound of confidence interval for harmonic mean
<code>r(Var_h)</code>	variance of $1/x_i$

Methods and formulas

See [Armitage, Berry, and Matthews \(2002\)](#) or [Snedecor and Cochran \(1989\)](#). For a history of the concept of the mean, see [Plackett \(1958\)](#).

When restricted to the same set of values (that is, to positive values), the arithmetic mean (\bar{x}) is greater than or equal to the geometric mean, which in turn is greater than or equal to the harmonic mean. Equality holds only if all values within a sample are equal to a positive constant.

The arithmetic mean and its confidence interval are identical to those provided by `ci`; see [\[R\] ci](#).

To compute the geometric mean, `ameans` first creates $u_j = \ln x_j$ for all positive x_j . The arithmetic mean of the u_j and its confidence interval are then computed as in `ci`. Let \bar{u} be the resulting mean, and let $[L, U]$ be the corresponding confidence interval. The geometric mean is then $\exp(\bar{u})$, and its confidence interval is $[\exp(L), \exp(U)]$.

The same procedure is followed for the harmonic mean, except that then $u_j = 1/x_j$. The harmonic mean is then $1/\bar{u}$, and its confidence interval is $[1/U, 1/L]$ if L is greater than zero. If L is not greater than zero, this confidence interval is not defined, and missing values are reported.

When weights are specified, `ameans` applies the weights to the transformed values, $u_j = \ln x_j$ and $u_j = 1/x_j$, respectively, when computing the geometric and harmonic means. For details on how the weights are used to compute the mean and variance of the u_j , see [\[R\] summarize](#). Without weights, the formula for the geometric mean reduces to

$$\exp\left\{\frac{1}{n} \sum_j \ln(x_j)\right\}$$

Without weights, the formula for the harmonic mean is

$$\frac{n}{\sum_j \frac{1}{x_j}}$$

Acknowledgments

This improved version of `ameans` is based on the `gmc_i` command ([Carlin, Vidmar, and Ramalheira 1998](#)) and was written by John Carlin of the Murdoch Children's Research Institute and the University of Melbourne; Suzanna Vidmar of the University of Melbourne; and Carlos Ramalheira of Coimbra University Hospital, Portugal.

References

- Armitage, P., G. Berry, and J. N. S. Matthews. 2002. *Statistical Methods in Medical Research*. 4th ed. Oxford: Blackwell.
- Carlin, J. B., S. Vidmar, and C. Ramalheira. 1998. [sg75: Geometric means and confidence intervals](#). *Stata Technical Bulletin* 41: 23–25. Reprinted in *Stata Technical Bulletin Reprints*, vol. 7, pp. 197–199. College Station, TX: Stata Press.
- Keynes, J. M. 1911. The principal averages and the laws of error which lead to them. *Journal of the Royal Statistical Society* 74: 322–331.
- Plackett, R. L. 1958. Studies in the history of probability and statistics: VII. The principle of the arithmetic mean. *Biometrika* 45: 130–135.
- Snedecor, G. W., and W. G. Cochran. 1989. *Statistical Methods*. 8th ed. Ames, IA: Iowa State University Press.
- Stigler, S. M. 1985. Arithmetic means. In Vol. 1 of *Encyclopedia of Statistical Sciences*, ed. S. Kotz and N. L. Johnson, 126–129. New York: Wiley.

Also see

[R] **ci** — Confidence intervals for means, proportions, and counts

[R] **mean** — Estimate means

[R] **summarize** — Summary statistics

[SVY] **svy estimation** — Estimation commands for survey data