

## Title

**halton()** — Generate a Halton or Hammersley sequence

## Syntax

```
real matrix halton(real scalar n, d [, real scalar burn [, real scalar method]])  
void          _halton(real matrix x, [, real scalar burn [, real scalar method]])
```

where inputs are

*n*: length of the sequence  
*d*: dimension of the sequence ( $\leq 20$ )  
*x*: an  $n \times d$  real matrix  
*burn*: starting index of the sequence, default 0  
*method*: type of sequence, 1=Halton (default), 2=Hammersley

## Description

This routine generates either a Halton, the default, or a Hammersley sequence of length  $n$  and dimension  $d \leq 20$ . These sequences have a more uniform scatter over the unit hypercube than sequences generated from pseudo-uniform random numbers.

## Remarks

The Halton sequence of dimension  $d$  is generated from the first  $d$  primes,  $p_k$ , so that on draw  $i$  we have  $\mathbf{h}_i = (r_{p_1}(i), r_{p_2}(i), \dots, r_{p_d}(i))$ , where

$$r_{p_k}(i) = \sum_{j=0}^q b_{jk}(i) p_k^{-j-1} \in (0, 1)$$

is the radical inverse function of  $i$  with base  $p_k$  so that  $\sum_{j=0}^q b_{jk}(i) p_k^j = i$  where  $p_k^q \leq i < p_k^{q+1}$  (Fan and Wang, 1994).

We demonstrate with base  $p_3 = 5$ ,  $i = 33$ , and we will generate  $r_5(33)$ . Here,  $q = 2$ ,  $b_{0,3}(33) = 3$ ,  $b_{1,5}(33) = 1$  and  $b_{2,5}(33) = 1$ , so that  $r_5(33) = \frac{3}{5} + \frac{1}{25} + \frac{1}{625}$ .

The Hammersley sequence uses an evenly spaced set of points with the first  $d-1$  components the Halton sequence,  $\mathbf{h}_i = (\frac{2i-1}{2n}, r_{p_1}(i), r_{p_2}(i), \dots, r_{p_{d-1}}(i))$ ,  $i = 1, \dots, n$ .

The discrepancy of a sequence, denoted  $D(n, d)$ , is a measure of its nonuniformity over the unit hypercube. For all rectangles on the unit hypercube  $(\mathbf{0}, \mathbf{t})$ ,  $\mathbf{t} = (t_1, \dots, t_d)$ ,  $t_j \leq 1$ , it measures the maximum of the absolute value of the difference between the ratio of the number of points in  $(\mathbf{0}, \mathbf{t})$  and the total number of points,  $nd$ , and the volume of  $(\mathbf{0}, \mathbf{t})$ . See Section 1.4 of Fang and Wang (1994) for a more rigorous definition. The discrepancy of the uniform pseudo-random sequence has a convergence rate that is independent of  $d$ ,  $D_{uni}(n) \leq K \sqrt{\frac{\log \log n}{n}}$  for some  $K > 0$ . For the Halton sequence we have a faster rate of convergence,  $D_{hal}(n, d) \leq \frac{K(d)(\log n)^d}{n}$ , while the Hammersley sequence has an improvement over the Halton,  $D_{ham}(n, d) \leq \frac{K(d)(\log n)^{d-1}}{n}$ .

## Conformability

`halton(n, d, burn, method)`

input:

`n`: 1 x 1  
`d`: 1 x 1  
`burn`: 1 x 1  
`method`: 1 x 1

output:

`result`:  $n \times d$

`_halton(x, burn, method)`

input:

`x`:  $n \times d$   
`burn`: 1 x 1  
`method`: 1 x 1

output:

`x`:  $n \times d$

## Diagnostics

`halton(n, d)` returns an  $n \times d$  matrix containing  $d$  sequences of length  $n$ . The first row of the matrix will have the sequence at index `burn`.

`_halton(x)` modifies the  $n \times d$  matrix `x` so that it contains  $d$  sequences of length  $n$ .

The maximum dimension,  $d$ , is 20.

## References

Fang, K.T. and Y. Wang 1994. *Number-Theoretic Methods in Statistics* London:Chapman & Hall.