

## example 2 — Mixture of Poisson regression models

[Description](#)[Remarks and examples](#)[References](#)[Also see](#)

## Description

In this example, we demonstrate how to fit a two-component mixture of Poisson regressions models. We also use `estat lmean` to estimate marginal predicted counts and `estat lprob` to estimate the proportion of individuals in each class.

## Remarks and examples

[stata.com](#)

We are interested in fitting a Poisson regression to model the annual number of doctor visits. We hypothesize that there are two distinct groups or classes in the population that differ in their healthcare utilization—frequent users and infrequent users—and we believe that the model may differ across these two groups.

We do not have any information that tells us which individuals in our sample belong to which group. With FMM, we can specify two latent classes in our model to identify these groups. To account for differences between the latent classes, we include predictor variables in our model to fit potentially different Poisson distributions for each class.

Here we replicate the finite mixture Poisson regression example from [\[SEM\] example 53g](#). We use the following data:

```
. use http://www.stata-press.com/data/r15/gsem_mixture
(U.S. Medical Expenditure Panel Survey (2003))
. describe
Contains data from http://www.stata-press.com/data/r15/gsem_mixture.dta
  obs:      3,677          U.S. Medical Expenditure Panel
                   Survey (2003)
  vars:      12           26 Jan 2017 08:46
  size:     62,509       (_dta has notes)
```

variable name	storage type	display format	value label	variable label
drvisits	int	%9.0g		number of doctor visits
private	byte	%8.0g		has private supplementary insurance
medicaid	byte	%8.0g		has Medicaid public insurance
age	byte	%8.0g		age in years
educ	byte	%8.0g		years of education
actlim	byte	%8.0g		has activity limitations
chronic	byte	%8.0g		number of chronic conditions
income	float	%9.0g		income in \$1,000s
offer	byte	%8.0g		employer offers insurance
hpvisits	int	%8.0g		number of visits to health professionals other than doctors
female	byte	%8.0g		female
phylim	byte	%8.0g		has physical limitation

Sorted by:

## 2 example 2 — Mixture of Poisson regression models

- ```
. notes
```
- \_dta:
1. Data on annual number of doctor visits for individuals age 65 and older from the U.S. Medical Expenditure Panel Survey for 2003.
  2. Data is analyzed in Cameron, A. C. and P. K. Trivedi, 2010, *Microeconometrics Using Stata*, Rev. Ed., College Station, TX: Stata Press.
  3. Additional information on finite mixture models for count data and a similar example are found in Deb, P. and P. K. Trivedi, 1997, Demand for medical care by the elderly: A finite mixture approach, *Journal of Applied Econometrics*, vol. 12, 313--336.

Following [Cameron and Trivedi \(2010\)](#), we fit an FMM with a Poisson regression component for each latent class. We model the number of doctor visits as a function of whether an individual has private supplementary insurance, whether he or she has Medicaid, age, age squared, education level, whether he or she has activity limitations, and the number of chronic conditions.

We add the `startvalues(randomid, draws(5) seed(15))` option to specify that five random draws are taken when computing starting values. The class assignment is selected from the draw that has the best log likelihood after the EM iterations. When fitting FMMS, taking multiple draws of random starting values can help to prevent convergence at a local maximum rather than the global maximum. `fmm` provides a variety of options for obtaining starting values; see [\[FMM\] fmm](#) for more information on starting values.

```
. fmm 2, startvalues(randomid, draws(5) seed(15)):
> poisson drvisits private medicaid c.age#c.age educ actlim chronic
      (iteration log omitted)
Finite mixture model                Number of obs      =      3,677
Log likelihood = -11502.686
```

|         | Coef.          | Std. Err. | z     | P> z  | [95% Conf. Interval] |          |
|---------|----------------|-----------|-------|-------|----------------------|----------|
| 1.Class | (base outcome) |           |       |       |                      |          |
| 2.Class |                |           |       |       |                      |          |
| _cons   | .877227        | .0494614  | 17.74 | 0.000 | .7802845             | .9741696 |

```
Class      : 1
Response   : drvisits
Model      : poisson
```

|             | Coef.     | Std. Err. | z     | P> z  | [95% Conf. Interval] |           |
|-------------|-----------|-----------|-------|-------|----------------------|-----------|
| drvisits    |           |           |       |       |                      |           |
| private     | .138229   | .0247626  | 5.58  | 0.000 | .0896951             | .1867629  |
| medicaid    | .1269723  | .0341525  | 3.72  | 0.000 | .0600345             | .19391    |
| age         | .2628874  | .0466774  | 5.63  | 0.000 | .1714014             | .3543735  |
| c.age#c.age | -.0017418 | .0003108  | -5.60 | 0.000 | -.002351             | -.0011326 |
| educ        | .0241679  | .0030705  | 7.87  | 0.000 | .0181499             | .030186   |
| actlim      | .1831598  | .0238817  | 7.67  | 0.000 | .1363525             | .2299671  |
| chronic     | .1970511  | .0088783  | 22.19 | 0.000 | .17965               | .2144523  |
| _cons       | -8.051256 | 1.741677  | -4.62 | 0.000 | -11.46488            | -4.637632 |

```

Class      : 2
Response   : drvisits
Model      : poisson

```

|             | Coef.     | Std. Err. | z     | P> z  | [95% Conf. Interval] |           |
|-------------|-----------|-----------|-------|-------|----------------------|-----------|
| drvisits    |           |           |       |       |                      |           |
| private     | .2077415  | .0306353  | 6.78  | 0.000 | .1476974             | .2677856  |
| medicaid    | .1071618  | .0407211  | 2.63  | 0.008 | .02735               | .1869736  |
| age         | .3798087  | .0562035  | 6.76  | 0.000 | .269652              | .4899655  |
| c.age#c.age | -.0024869 | .0003736  | -6.66 | 0.000 | -.0032191            | -.0017547 |
| educ        | .029099   | .003972   | 7.33  | 0.000 | .021314              | .0368841  |
| actlim      | .1244235  | .0310547  | 4.01  | 0.000 | .0635574             | .1852895  |
| chronic     | .3191166  | .0089757  | 35.55 | 0.000 | .3015247             | .3367086  |
| _cons       | -14.25713 | 2.101964  | -6.78 | 0.000 | -18.37691            | -10.13736 |

The first table in the output provides the estimated coefficients in the multinomial logit model for the latent class probabilities. The next two tables are the results for the Poisson regression models for the first and second classes. The estimated coefficients from these tables are interpreted just as you would coefficients from `poisson`; see [R] [poisson](#).

To better understand these classes, we use `estat lcmean` to estimate the marginal predicted counts (means) for each class.

```

. estat lcmean
Latent class marginal means          Number of obs   =      3,677

```

|          | Delta-method |           |       |       |                      |
|----------|--------------|-----------|-------|-------|----------------------|
|          | Margin       | Std. Err. | z     | P> z  | [95% Conf. Interval] |
| 1        |              |           |       |       |                      |
| drvisits | 13.95943     | .1767506  | 78.98 | 0.000 | 13.613 14.30585      |
| 2        |              |           |       |       |                      |
| drvisits | 3.801692     | .0587685  | 64.69 | 0.000 | 3.686508 3.916876    |

We see that class 1 represents those who visit the doctor frequently and class 2 represents those who visit the doctor less frequently. We can use `estat lcprob` to estimate the proportion of individuals in each class.

```

. estat lcprob
Latent class marginal probabilities    Number of obs   =      3,677


```

| Class | Delta-method |           |                      |
|-------|--------------|-----------|----------------------|
|       | Margin       | Std. Err. | [95% Conf. Interval] |
| 1     | .2937527     | .0102614  | .2740502 .3142586    |
| 2     | .7062473     | .0102614  | .6857414 .7259498    |

We find that about 29% of the population is in the group that visits the doctor frequently (class 1) and about 71% is in the group that visits the doctor less frequently (class 2).

We can visually compare the resulting distributions of the means by plotting the predicted number of doctor visits.

```
. predict mu*  
(option mu assumed)  
. twoway histogram mu1, width(1) color(navy) fcolor(%25) lcolor(%25) ||  
> histogram mu2, width(1) color(maroon) fcolor(%25) lcolor(%25)  
> legend(off) title("Predicted doctor visits for two groups")
```



We can clearly see the two groups. The frequent user group exhibits more variability, which is expected in a Poisson process where the variance is equal to the mean.

## References

- Cameron, A. C., and P. K. Trivedi. 2010. *Microeconometrics Using Stata*. Rev. ed. College Station, TX: Stata Press.
- Deb, P., and P. K. Trivedi. 1997. Demand for medical care by the elderly: A finite mixture approach. *Journal of Applied Econometrics* 12: 313–336.

## Also see

- [FMM] [fmm intro](#) — Introduction to finite mixture models
- [FMM] [fmm: poisson](#) — Finite mixtures of Poisson regression models
- [FMM] [estat lcmean](#) — Latent class marginal means
- [FMM] [estat lcprob](#) — Latent class marginal probabilities
- [SEM] [example 53g](#) — Finite mixture Poisson regression
- [SEM] [example 54g](#) — Finite mixture Poisson regression, multiple responses
- [SEM] [gsem](#) — Generalized structural equation model estimation command