

**hausman** — Hausman specification test[Description](#)[Quick start](#)[Menu](#)[Syntax](#)[Options](#)[Remarks and examples](#)[Stored results](#)[Methods and formulas](#)[Acknowledgment](#)[References](#)[Also see](#)

## Description

`hausman` performs Hausman's (1978) specification test.

## Quick start

Hausman test for stored models `consistent` and `efficient`

```
hausman consistent efficient
```

As above, but compare fixed-effects and random-effects linear regression models

```
hausman fixed random, sigmamore
```

Endogeneity test after `ivprobit` and `probit` with estimates stored in `iv` and `noiv`

```
hausman iv noiv, equations(1:1)
```

Test of independence of irrelevant alternatives for model with all alternatives `all` and model with omitted alternative `omitted`

```
hausman omitted all, alleqs constant
```

## Menu

Statistics > Postestimation

## Syntax

```
hausman name-consistent [name-efficient] [, options]
```

<i>options</i>	Description
Main	
<code>constant</code>	include estimated intercepts in comparison; default is to exclude
<code>alleqs</code>	use all equations to perform test; default is first equation only
<code>skipeqs(<i>eqlist</i>)</code>	skip specified equations when performing test
<code>equations(<i>matchlist</i>)</code>	associate/compare the specified (by number) pairs of equations
<code>force</code>	force performance of test, even though assumptions are not met
<code>df(#)</code>	use # degrees of freedom
<code>sigmamore</code>	base both (co)variance matrices on disturbance variance estimate from efficient estimator
<code>sigmaless</code>	base both (co)variance matrices on disturbance variance estimate from consistent estimator
Advanced	
<code>tconsistent(<i>string</i>)</code>	consistent estimator column header
<code>tefficient(<i>string</i>)</code>	efficient estimator column header

where *name-consistent* and *name-efficient* are names under which estimation results were stored via `estimates store`; see [R] [estimates store](#).

A period (.) may be used to refer to the last estimation results, even if these were not already stored.

Not specifying *name-efficient* is equivalent to specifying the last estimation results as “.”.

## Options

### Main

`constant` specifies that the estimated intercept(s) be included in the model comparison; by default, they are excluded. The default behavior is appropriate for models in which the constant does not have a common interpretation across the two models.

`alleqs` specifies that all the equations in the models be used to perform the Hausman test; by default, only the first equation is used.

`skipeqs(eqlist)` specifies in *eqlist* the names of equations to be excluded from the test. Equation numbers are not allowed in this context, because the equation names, along with the variable names, are used to identify common coefficients.

`equations(matchlist)` specifies, by number, the pairs of equations that are to be compared.

The *matchlist* in `equations()` should follow the syntax

$$\#_c:\#_e \left[ , \#_c:\#_e \left[ , \dots \right] \right]$$

where  $\#_c$  ( $\#_e$ ) is an equation number of the always-consistent (efficient under  $H_0$ ) estimator. For instance, `equations(1:1)`, `equations(1:1, 2:2)`, or `equations(1:2)`.

If `equations()` is not specified, then equations are matched on equation names.

`equations()` handles the situation in which one estimator uses equation names and the other does not. For instance, `equations(1:2)` means that equation 1 of the always-consistent estimator is to be tested against equation 2 of the efficient estimator. `equations(1:1, 2:2)` means that equation 1 is to be tested against equation 1 and that equation 2 is to be tested against equation 2. If `equations()` is specified, the `alleqs` and `skipeqs` options are ignored.

`force` specifies that the Hausman test be performed, even though the assumptions of the Hausman test seem not to be met, for example, because the estimators were `pweighted` or the data were clustered.

`df(#)` specifies the degrees of freedom for the Hausman test. The default is the matrix rank of the variance of the difference between the coefficients of the two estimators.

`sigmamore` and `sigmaless` specify that the two covariance matrices used in the test be based on a common estimate of disturbance variance ( $\sigma^2$ ).

`sigmamore` specifies that the covariance matrices be based on the estimated disturbance variance from the efficient estimator. This option provides a proper estimate of the contrast variance for so-called tests of exogeneity and overidentification in instrumental-variables regression.

`sigmaless` specifies that the covariance matrices be based on the estimated disturbance variance from the consistent estimator.

These options can be specified only when both estimators store `e(sigma)` or `e(rmse)`, or with the `xtreg` command. `e(sigma_e)` is stored after the `xtreg` command with the `fe` or `mle` option. `e(rmse)` is stored after the `xtreg` command with the `re` option.

`sigmamore` or `sigmaless` are recommended when comparing fixed-effects and random-effects linear regression because they are much less likely to produce a non-positive-definite-differenced covariance matrix (although the tests are asymptotically equivalent whether or not one of the options is specified).

#### Advanced

`tconsistent(string)` and `tefficient(string)` are formatting options. They allow you to specify the headers of the columns of coefficients that default to the names of the models. These options will be of interest primarily to programmers.

## Remarks and examples

[stata.com](http://www.stata.com)

`hausman` is a general implementation of Hausman's (1978) specification test, which compares an estimator  $\hat{\theta}_1$  that is known to be consistent with an estimator  $\hat{\theta}_2$  that is efficient under the assumption being tested. The null hypothesis is that the estimator  $\hat{\theta}_2$  is indeed an efficient (and consistent) estimator of the true parameters. If this is the case, there should be no systematic difference between the two estimators. If there exists a systematic difference in the estimates, you have reason to doubt the assumptions on which the efficient estimator is based.

The assumption of efficiency is violated if the estimator is `pweighted` or the data are clustered, so `hausman` cannot be used. The test can be forced by specifying the `force` option with `hausman`. For an alternative to using `hausman` in these cases, see [R] [suest](#).

To use `hausman`, you

```
. (compute the always-consistent estimator)
. estimates store name-consistent
. (compute the estimator that is efficient under H_0)
. hausman name-consistent .
```

Alternatively, you can turn this around:

```
. (compute the estimator that is efficient under  $H_0$ )
. estimates store name-efficient
. (fit the less-efficient model)
. (compute the always-consistent estimator)
. hausman . name-efficient
```

You can, of course, also compute and store both the always-consistent and efficient-under- $H_0$  estimators and perform the Hausman test with

```
. hausman name-consistent name-efficient
```

### ► Example 1

We are studying the factors that affect the wages of young women in the United States between 1970 and 1988, and we have a panel-data sample of individual women over that time span.

```
. use https://www.stata-press.com/data/r16/nlswork4
(National Longitudinal Survey. Young Women 14-26 years of age in 1968)
. describe
Contains data from https://www.stata-press.com/data/r16/nlswork4.dta
  obs:      28,534      National Longitudinal Survey.
                        Young Women 14-26 years of age
                        in 1968
  vars:      6         29 Jan 2018 16:35
```

---

variable name	storage type	display format	value label	variable label
idcode	int	%8.0g		NLS ID
year	byte	%8.0g		interview year
age	byte	%8.0g		age in current year
msp	byte	%8.0g		1 if married, spouse present
ttl_exp	float	%9.0g		total work experience
ln_wage	float	%9.0g		ln(wage/GNP deflator)

---

```
Sorted by: idcode year
```

We believe that a random-effects specification is appropriate for individual-level effects in our model. We fit a fixed-effects model that will capture all temporally constant individual-level effects.

```
. xtreg ln_wage age msp ttl_exp, fe
```

```
Fixed-effects (within) regression      Number of obs   =   28,494
Group variable: idcode                 Number of groups =    4,710
R-sq:                                  Obs per group:
    within = 0.1373                      min =           1
    between = 0.2571                      avg =           6.0
    overall = 0.1800                      max =           15
                                         F(3,23781)     =   1262.01
                                         Prob > F       =    0.0000
```

```
corr(u_i, Xb) = 0.1476
```

ln_wage	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	-.005485	.000837	-6.55	0.000	-.0071256	-.0038443
msp	.0033427	.0054868	0.61	0.542	-.0074118	.0140971
ttl_exp	.0383604	.0012416	30.90	0.000	.0359268	.0407941
_cons	1.593953	.0177538	89.78	0.000	1.559154	1.628752
sigma_u	.37674223					
sigma_e	.29751014					
rho	.61591044	(fraction of variance due to u_i)				

```
F test that all u_i=0: F(4709, 23781) = 7.76          Prob > F = 0.0000
```

We assume that this model is consistent for the true parameters and store the results by using `estimates store` under a name, `fixed`:

```
. estimates store fixed
```

Now we fit a random-effects model as a fully efficient specification of the individual effects under the assumption that they are random and follow a normal distribution. We then compare these estimates with the previously stored results by using the `hausman` command.

```
. xtreg ln_wage age msp ttl_exp, re
```

```
Random-effects GLS regression      Number of obs   =   28,494
Group variable: idcode             Number of groups =    4,710
R-sq:                               Obs per group:
    within = 0.1373                      min =           1
    between = 0.2552                      avg =           6.0
    overall = 0.1797                      max =           15
                                         Wald chi2(3)   =   5100.33
                                         Prob > chi2    =    0.0000
```

```
corr(u_i, X) = 0 (assumed)
```

ln_wage	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
age	-.0069749	.0006882	-10.13	0.000	-.0083238	-.0056259
msp	.0046594	.0051012	0.91	0.361	-.0053387	.0146575
ttl_exp	.0429635	.0010169	42.25	0.000	.0409704	.0449567
_cons	1.609916	.0159176	101.14	0.000	1.578718	1.641114
sigma_u	.32648519					
sigma_e	.29751014					
rho	.54633481	(fraction of variance due to u_i)				

```
. hausman fixed ., sigmamore
```

	Coefficients		(b-B) Difference	sqrt(diag(V_b-V_B)) S.E.
	(b) fixed	(B) .		
age	-.005485	-.0069749	.0014899	.0004803
msp	.0033427	.0046594	-.0013167	.0020596
t1l_exp	.0383604	.0429635	-.0046031	.0007181

b = consistent under Ho and Ha; obtained from xtreg

B = inconsistent under Ha, efficient under Ho; obtained from xtreg

Test: Ho: difference in coefficients not systematic

$$\text{chi2}(3) = (b-B)'[(V_b-V_B)^{-1}](b-B)$$

$$= 260.40$$

$$\text{Prob}>\text{chi2} = 0.0000$$

Under the current specification, our initial hypothesis that the individual-level effects are adequately modeled by a random-effects model is resoundingly rejected. This result is based on the rest of our model specification, and random effects might be appropriate for some alternate model of wages. ◀

Jerry Allen Hausman (1946– ) is an American economist and econometrician. He was born in West Virginia and went on to study economics at Brown and Oxford. He joined the MIT faculty in 1972 and continues to teach there. He currently researches new goods and their effects on consumer welfare and its measurement in the Consumer Price Index along with regulation and competition in the telecommunications industry.

Hausman is best known for his many contributions to econometrics. In 1978, he published his now famous paper giving the Hausman specification test. The work remains one of the most widely cited econometrics papers. He has also done extensive work in applied microeconomics pertaining to governments role in the economy, including antitrust regulation, public finance, and taxation.

In 1980, Hausman received the Frisch Medal, a biennial award from the Econometric Society recognizing exceptional applied work, for his paper with David Wise on attrition bias. In 1985, he won the John Bates Clark Award from the American Economics Association, which is given for outstanding contributions to economics by an economist under 40 years of age. In 2012, the *Advances in Econometrics* book series devoted an entire volume to Hausman and his contributions to econometrics.

## ▷ Example 2

A stringent assumption of multinomial and conditional logit models is that outcome categories for the model have the property of independence of irrelevant alternatives (IIA). Stated simply, this assumption requires that the inclusion or exclusion of categories does not affect the relative risks associated with the regressors in the remaining categories.

One classic example of a situation in which this assumption would be violated involves the choice of transportation mode; see [McFadden \(1974\)](#). For simplicity, postulate a transportation model with the four possible outcomes: rides a train to work, takes a bus to work, drives the Ford to work, and drives the Chevrolet to work. Clearly, “drives the Ford” is a closer substitute to “drives the Chevrolet” than it is to “rides a train” (at least for most people). This means that excluding “drives the Ford” from the model could be expected to affect the relative risks of the remaining options and that the model would not obey the IIA assumption.

Using the data presented in [R] [mlogit](#), we will use a simplified model to test for IIA. The choice of insurance type among indemnity, prepaid, and uninsured is modeled as a function of age and gender. The indemnity category is allowed to be the base category, and the model including all three outcomes is fit. The results are then stored under the name `allcats`.

```
. use https://www.stata-press.com/data/r16/sydsdn3
(Health insurance data)
. mlogit insure age male
Iteration 0:  log likelihood = -555.85446
Iteration 1:  log likelihood = -551.32973
Iteration 2:  log likelihood = -551.32802
Iteration 3:  log likelihood = -551.32802

Multinomial logistic regression           Number of obs   =           615
                                           LR chi2(4)      =           9.05
                                           Prob > chi2     =          0.0598
Log likelihood = -551.32802              Pseudo R2       =          0.0081
```

insure	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
Indemnity (base outcome)						
Prepaid						
age	-.0100251	.0060181	-1.67	0.096	-.0218204	.0017702
male	.5095747	.1977893	2.58	0.010	.1219147	.8972346
_cons	.2633838	.2787575	0.94	0.345	-.2829708	.8097383
Uninsure						
age	-.0051925	.0113821	-0.46	0.648	-.0275011	.0171161
male	.4748547	.3618462	1.31	0.189	-.2343508	1.18406
_cons	-1.756843	.5309602	-3.31	0.001	-2.797506	-.7161803

```
. estimates store allcats
```

Under the IIA assumption, we would expect no systematic change in the coefficients if we excluded one of the outcomes from the model. (For an extensive discussion, see [Hausman and McFadden \[1984\]](#).) We reestimate the parameters, excluding the uninsured outcome, and perform a Hausman test against the fully efficient full model.

```
. mlogit insure age male if insure != "Uninsure":insure
Iteration 0:  log likelihood = -394.8693
Iteration 1:  log likelihood = -390.4871
Iteration 2:  log likelihood = -390.48643
Iteration 3:  log likelihood = -390.48643

Multinomial logistic regression           Number of obs   =           570
                                           LR chi2(2)      =           8.77
                                           Prob > chi2     =          0.0125
Log likelihood = -390.48643              Pseudo R2       =          0.0111
```

insure	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
Indemnity (base outcome)						
Prepaid						
age	-.0101521	.0060049	-1.69	0.091	-.0219214	.0016173
male	.5144003	.1981735	2.60	0.009	.1259874	.9028133
_cons	.2678043	.2775563	0.96	0.335	-.276196	.8118046

```
. hausman . allcats, alleqs constant
```

	Coefficients		(b-B) Difference	sqrt(diag(V_b-V_B)) S.E.
	(b)	(B) allcats		
age	-.0101521	-.0100251	-.0001269	.
male	.5144003	.5095747	.0048256	.0123338
_cons	.2678043	.2633838	.0044205	.

b = consistent under Ho and Ha; obtained from mlogit  
 B = inconsistent under Ha, efficient under Ho; obtained from mlogit

Test: Ho: difference in coefficients not systematic

chi2(3) = (b-B)'[(V\_b-V\_B)<sup>-1</sup>](b-B)  
 = 0.08  
 Prob>chi2 = 0.9944  
 (V\_b-V\_B is not positive definite)

The syntax of the if condition on the mlogit command simply identified the "Uninsured" category with the insure value label; see [U] 12.6.3 Value labels. On examining the output from hausman, we see that there is no evidence that the IIA assumption has been violated.

Because the Hausman test is a standardized comparison of model coefficients, using it with mlogit requires that the base outcome be the same in both competing models. In particular, if the most-frequent category (the default base outcome) is being removed to test for IIA, you must use the baseoutcome() option in mlogit to manually set the base outcome to something else. Or you can use the equation() option of the hausman command to align the equations of the two models.

Having the missing values for the square root of the diagonal of the covariance matrix of the differences is not comforting, but it is also not surprising. This covariance matrix is guaranteed to be positive definite only asymptotically (it is a consequence of the assumption that one of the estimators is efficient), and assurances are not made about the diagonal elements. Negative values along the diagonal are possible, and the fourth column of the table is provided mainly for descriptive use.

We can also perform the Hausman IIA test against the remaining alternative in the model:

```
. mlogit insure age male if insure != "Prepaid":insure
Iteration 0: log likelihood = -132.59913
Iteration 1: log likelihood = -131.78009
Iteration 2: log likelihood = -131.76808
Iteration 3: log likelihood = -131.76807

Multinomial logistic regression           Number of obs   =       338
                                           LR chi2(2)      =         1.66
                                           Prob > chi2     =       0.4356
                                           Pseudo R2      =       0.0063

Log likelihood = -131.76807
```

insure	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
Indemnity	(base outcome)					
Uninsure						
age	-.0041055	.0115807	-0.35	0.723	-.0268033	.0185923
male	.4591074	.3595663	1.28	0.202	-.2456296	1.163844
_cons	-1.801774	.5474476	-3.29	0.001	-2.874752	-.7287968



```
. hausman . allcats, alleqs constant
```

	Coefficients		(b-B) Difference	sqrt(diag(V_b-V_B)) S.E.
	(b)	(B) allcats		
age	-.0041055	-.0051925	.001087	.0021355
male	.4591074	.4748547	-.0157473	.
_cons	-1.801774	-1.756843	-.0449311	.1333421

b = consistent under Ho and Ha; obtained from mlogit  
 B = inconsistent under Ha, efficient under Ho; obtained from mlogit

Test: Ho: difference in coefficients not systematic

```
chi2(3) = (b-B)'[(V_b-V_B)^(-1)](b-B)
         = -0.18      chi2<0 ==> model fitted on these
                        data fails to meet the asymptotic
                        assumptions of the Hausman test;
                        see suest for a generalized test
```

Here the  $\chi^2$  statistic is actually negative. We might interpret this result as strong evidence that we cannot reject the null hypothesis. Such a result is not an unusual outcome for the Hausman test, particularly when the sample is relatively small—there are only 45 uninsured individuals in this dataset.

Are we surprised by the results of the Hausman test in this example? Not really. Judging from the  $z$  statistics on the original multinomial logit model, we were struggling to identify any structure in the data with the current specification. Even when we were willing to assume IIA and computed the efficient estimator under this assumption, few of the effects could be identified as statistically different from those on the base category. Trying to base a Hausman test on a contrast (difference) between two poor estimates is just asking too much of the existing data.

◀

In [example 2](#), we encountered a case in which the Hausman was not well defined. Unfortunately, in our experience this happens fairly often. Stata provides an alternative to the Hausman test that overcomes this problem through an alternative estimator of the variance of the difference between the two estimators. This other estimator is guaranteed to be positive semidefinite. This alternative estimator also allows a widening of the scope of problems to which Hausman-type tests can be applied by relaxing the assumption that one of the estimators is efficient. For instance, you can perform Hausman-type tests to clustered observations and survey estimators. See [\[R\] suest](#) for details.

## Stored results

hausman stores the following in `r()`:

Scalars

```
r(chi2)       $\chi^2$ 
r(df)       degrees of freedom for the statistic
r(p)         $p$ -value for the  $\chi^2$ 
r(rank)     rank of  $(V\_b-V\_B)^{-1}$ 
```

## Methods and formulas

The Hausman statistic is distributed as  $\chi^2$  and is computed as

$$H = (\beta_c - \beta_e)'(V_c - V_e)^{-1}(\beta_c - \beta_e)$$

where

$\beta_c$	is the coefficient vector from the consistent estimator
$\beta_e$	is the coefficient vector from the efficient estimator
$V_c$	is the covariance matrix of the consistent estimator
$V_e$	is the covariance matrix of the efficient estimator

When the difference in the variance matrices is not positive definite, a Moore–Penrose generalized inverse is used. As noted in [Gourieroux and Monfort \(1995, 125–128\)](#), the choice of generalized inverse is not important asymptotically.

The number of degrees of freedom for the statistic is the rank of the difference in the variance matrices. When the difference is positive definite, this is the number of common coefficients in the models being compared.

## Acknowledgment

Portions of `hausman` are based on an early implementation by Jeroen Weesie of the Department of Sociology at Utrecht University, The Netherlands.

## References

- Baltagi, B. H. 2011. *Econometrics*. 5th ed. Berlin: Springer.
- Gourieroux, C. S., and A. Monfort. 1995. *Statistics and Econometric Models, Vol 2: Testing, Confidence Regions, Model Selection, and Asymptotic Theory*. Trans. Q. Vuong. Cambridge: Cambridge University Press.
- Hausman, J. A. 1978. Specification tests in econometrics. *Econometrica* 46: 1251–1271.
- Hausman, J. A., and D. L. McFadden. 1984. Specification tests for the multinomial logit model. *Econometrica* 52: 1219–1240.
- McFadden, D. L. 1974. Measurement of urban travel demand. *Journal of Public Economics* 3: 303–328.

## Also see

- [R] [lrtest](#) — Likelihood-ratio test after estimation
- [R] [suest](#) — Seemingly unrelated estimation
- [R] [test](#) — Test linear hypotheses after estimation
- [XT] [xtreg](#) — Fixed-, between-, and random-effects and population-averaged linear models