

example 13 — Equation-level Wald test
[Description](#)[Remarks and examples](#)[Also see](#)

Description

We demonstrate `estat eqtest`. See [\[SEM\] intro 7](#) and [\[SEM\] estat eqtest](#).

This example picks up where [\[SEM\] example 12](#) left off:

```
. use http://www.stata-press.com/data/r15/auto
. sem (price <- foreign mpg displacement)          ///
      (weight <- foreign length),                ///
      cov(e.price*e.weight)
```

Remarks and examples

stata.com

We have fit a two-equation model with equations for endogenous variables `price` and `weight`. There happen to be two equations, the model happens to be a seemingly unrelated regression, and the endogenous variables happen to be observed, but none of that is important right now.

`estat eqtest` displays equation-by-equation Wald tests that all coefficients excluding the intercepts are 0.

```
. estat eqtest
Wald tests for equations
```

	chi2	df	p
observed			
price	36.43	3	0.0000
weight	633.34	2	0.0000

Note:

1. The null hypothesis for this test is that the coefficients other than the intercepts are 0. We can reject that null hypothesis for each equation.

Also see

[\[SEM\] example 12](#) — Seemingly unrelated regression

[\[SEM\] intro 7](#) — Postestimation tests and predictions

[\[SEM\] estat eqtest](#) — Equation-level test that all coefficients are zero