

xtologit — Random-effects ordered logistic models

Description	Quick start	Menu	Syntax
Options	Remarks and examples	Stored results	Methods and formulas
References	Also see		

Description

`xtologit` fits random-effects ordered logistic models. The actual values taken on by the dependent variable are irrelevant, although larger values are assumed to correspond to “higher” outcomes. The conditional distribution of the dependent variable given the random effects is assumed to be multinomial with success probability determined by the logistic cumulative distribution function.

Quick start

Random-effects ordered logistic model of y as a function of x using `xtset` data

```
xtologit y x
```

Add [indicators](#) for levels of categorical variable a

```
xtologit y x i.a
```

As above, but report odds ratios

```
xtologit y x i.a, or
```

With cluster-robust standard errors for panels nested within `cvar`

```
xtologit y x i.a, vce(cluster cvar)
```

Menu

Statistics > Longitudinal/panel data > Ordinal outcomes > Logistic regression (RE)

Syntax

```
xtlogit depvar [indepvars] [if] [in] [weight] [, options]
```

<i>options</i>	Description
Model	
<code>offset(<i>varname</i>)</code>	include <i>varname</i> in model with coefficient constrained to 1
<code>constraints(<i>constraints</i>)</code>	apply specified linear constraints
<code>collinear</code>	keep collinear variables
SE/Robust	
<code>vce(<i>vcetype</i>)</code>	<i>vcetype</i> may be <code>oim</code> , <code>robust</code> , <code>cluster <i>clustvar</i></code> , <code>bootstrap</code> , or <code>jackknife</code>
Reporting	
<code>level(#)</code>	set confidence level; default is <code>level(95)</code>
<code>or</code>	report odds ratios
<code>lrmodel</code>	perform the likelihood-ratio model test instead of the default Wald test
<code>nocnsreport</code>	do not display constraints
<code>disply_<options></code>	control columns and column formats, row spacing, line width, display of omitted variables and base and empty cells, and factor-variable labeling
Integration	
<code>intmethod(<i>intmethod</i>)</code>	integration method; <i>intmethod</i> may be <code>mvaghermite</code> (the default) or <code>ghermite</code>
<code>intpoints(#)</code>	use # quadrature points; default is <code>intpoints(12)</code>
Maximization	
<code>maximize_<options></code>	control the maximization process; seldom used
<code>startgrid(<i>numlist</i>)</code>	improve starting value of the random-intercept parameter by performing a grid search
<code>nodisplay</code>	suppress display of header and coefficients
<code>coeflegend</code>	display legend instead of statistics

A panel variable must be specified; see [XT] [xtset](#).

indepvars may contain factor variables; see [U] [11.4.3 Factor variables](#).

depvar and *indepvars* may contain time-series operators; see [U] [11.4.4 Time-series varlists](#).

`by`, `fp`, and `statsby` are allowed; see [U] [11.1.10 Prefix commands](#).

`fweights`, `iweights`, and `pweights` are allowed; see [U] [11.1.6 weight](#).

`startgrid()`, `nodisplay`, and `coeflegend` do not appear in the dialog box.

See [U] [20 Estimation and postestimation commands](#) for more capabilities of estimation commands.

Options

Model

`offset(varname)`, `constraints(constraints)`, `collinear`; see [R] [estimation options](#).

SE/Robust

`vce(vcetype)` specifies the type of standard error reported, which includes types that are derived from asymptotic theory (`oim`), that are robust to some kinds of misspecification (`robust`), that allow for intragroup correlation (`cluster clustvar`), and that use bootstrap or jackknife methods (`bootstrap`, `jackknife`); see [XT] [vce_options](#).

Specifying `vce(robust)` is equivalent to specifying `vce(cluster panelvar)`; see [xtologit and the robust VCE estimator](#) in *Methods and formulas*.

Reporting

`level(#)`; see [R] [estimation options](#).

`or` reports the estimated coefficients transformed to odds ratios, that is, e^b rather than b . Standard errors and confidence intervals are similarly transformed. This option affects how results are displayed, not how they are estimated. `or` may be specified at estimation or when replaying previously estimated results.

`lrmodel`, `nocnsreport`; see [R] [estimation options](#).

`display_options`: `nocl`, `nopvalues`, `noomitted`, `vsquish`, `noemptycells`, `baselevels`, `allbaselevels`, `nofvlabel`, `fvwrap(#)`, `fvwrapon(style)`, `cformat(%fmt)`, `pformat(%fmt)`, `sformat(%fmt)`, and `nolstretch`; see [R] [estimation options](#).

Integration

`intmethod(intmethod)`, `intpoints(#)`; see [R] [estimation options](#).

Maximization

`maximize_options`: `difficult`, `technique(algorithm_spec)`, `iterate(#)`, `[no]log`, `trace`, `gradient`, `showstep`, `hessian`, `showtolerance`, `tolerance(#)`, `ltolerance(#)`, `nrtolerance(#)`, `nonrtolerance`, and `from(init_specs)`; see [R] [maximize](#). These options are seldom used.

The following options are available with `xtologit` but are not shown in the dialog box:

`startgrid(numlist)` performs a grid search to improve the starting value of the random-intercept parameter. No grid search is performed by default unless the starting value is found to not be feasible; in this case, `xtologit` runs `startgrid(0.1 1 10)` and chooses the value that works best. You may already be using a default form of `startgrid()` without knowing it. If you see `xtologit` displaying Grid node 1, Grid node 2, ... following Grid node 0 in the iteration log, that is `xtologit` doing a default search because the original starting value was not feasible.

`nodisplay` is for programmers. It suppresses the display of the header and the coefficients.

`coeflegend`; see [R] [estimation options](#).

Remarks and examples

Remarks are presented under the following headings:

[Overview](#)

[Video example](#)

Overview

`xtologit` fits random-effects ordered logistic models. Ordered logistic models are used to estimate relationships between an ordinal dependent variable and a set of independent variables. An *ordinal* variable is a variable that is categorical and ordered, for instance, “poor”, “good”, and “excellent”, which might indicate a person’s current health status or the repair record of a car. If there are only two outcomes, see [XT] [xtlogit](#), [XT] [xtprobit](#), and [XT] [xtcloglog](#). This entry is concerned only with more than two outcomes.

▷ Example 1

We use the data from the “Television, School, and Family Smoking Prevention and Cessation Project” (Flay et al. 1988; Rabe-Hesketh and Skrondal 2012, chap. 11), where schools were randomly assigned into one of four groups defined by two treatment variables. Students within each school are nested in classes, and classes are nested in schools. In this example, we ignore the variability of classes within schools; see [example 2](#) of [ME] [meologit](#) for a model that incorporates the additional class-level variance component. The dependent variable is the tobacco and health knowledge score (`thk`) collapsed into four ordered categories. We regress the outcome on the treatment variables and their interaction and control for the pretreatment score.

```

. use http://www.stata-press.com/data/r15/tvsfpors
. xtset school
      panel variable:  school (unbalanced)
. xtologit thk prethk cc##tv
Fitting comparison model:
Iteration 0:  log likelihood = -2212.775
Iteration 1:  log likelihood = -2125.509
Iteration 2:  log likelihood = -2125.1034
Iteration 3:  log likelihood = -2125.1032
Refining starting values:
Grid node 0:  log likelihood = -2136.2426
Fitting full model:
Iteration 0:  log likelihood = -2136.2426 (not concave)
Iteration 1:  log likelihood = -2120.2577
Iteration 2:  log likelihood = -2119.7574
Iteration 3:  log likelihood = -2119.7428
Iteration 4:  log likelihood = -2119.7428
Random-effects ordered logistic regression
Group variable:  school
Random effects u_i ~ Gaussian
Number of obs      =      1,600
Number of groups   =         28
Obs per group:
      min =         18
      avg =        57.1
      max =        137
Integration method: mvaghermite
Integration pts.   =         12
Wald chi2(4)      =        128.06
Prob > chi2       =         0.0000
Log likelihood = -2119.7428

```

thk	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
prethk	.4032892	.03886	10.38	0.000	.327125	.4794534
1.cc	.9237904	.204074	4.53	0.000	.5238127	1.323768
1.tv	.2749937	.1977424	1.39	0.164	-.1125744	.6625618
cc##tv						
1 1	-.4659256	.2845963	-1.64	0.102	-1.023724	.0918728
/cut1	-.0884493	.1641062			-.4100916	.233193
/cut2	1.153364	.165616			.8287625	1.477965
/cut3	2.33195	.1734199			1.992053	2.671846
/sigma2_u	.0735112	.0383106			.0264695	.2041551

```
LR test vs. ologit model:  chibar2(01) = 10.72      Prob >= chibar2 = 0.0005
```

The estimation table reports the parameter estimates, the estimated cutpoints ($\kappa_1, \kappa_2, \kappa_3$), and the estimated panel-level variance component labeled `sigma2_u`. The parameter estimates can be interpreted just as the output from a standard ordered logistic regression would be interpreted; see [R] [ologit](#). For example, we find that students with higher preintervention scores tend to have higher postintervention scores.

Underneath the parameter estimates and the cutpoints, the table shows the estimated variance component. The estimate of σ_u^2 is 0.074 with standard error 0.038. The reported likelihood-ratio test shows that there is enough variability between schools to favor a random-effects ordered logistic regression over a standard ordered logistic regression.

□ Technical note

The random-effects model is calculated using quadrature, which is an approximation whose accuracy depends partially on the number of integration points used. We can use the `quadchk` command to see if changing the number of integration points affects the results. If the results change, the quadrature approximation is not accurate given the number of integration points. Try increasing the number of integration points using the `intpoints()` option and run `quadchk` again. Do not attempt to interpret the results of estimates when the coefficients reported by `quadchk` differ substantially. See [\[XT\] quadchk](#) for details and [\[XT\] xtprobit](#) for an [example](#).

Because the `xtlogit` likelihood function is calculated by Gauss–Hermite quadrature, on large problems the computations can be slow. Computation time is roughly proportional to the number of points used for the quadrature.

□

Video example

[Ordered logistic and probit for panel data](#)

Stored results

`xtlogit` stores the following in `e()`:

Scalars

<code>e(N)</code>	number of observations
<code>e(N_g)</code>	number of groups
<code>e(k)</code>	number of parameters
<code>e(k_aux)</code>	number of auxiliary parameters
<code>e(k_eq)</code>	number of equations in <code>e(b)</code>
<code>e(k_eq_model)</code>	number of equations in overall model test
<code>e(k_dv)</code>	number of dependent variables
<code>e(k_cat)</code>	number of categories
<code>e(df_m)</code>	model degrees of freedom
<code>e(ll)</code>	log likelihood
<code>e(ll_0)</code>	log likelihood, constant-only model
<code>e(ll_c)</code>	log likelihood, comparison model
<code>e(chi2)</code>	χ^2
<code>e(chi2_c)</code>	χ^2 for comparison test
<code>e(N_clust)</code>	number of clusters
<code>e(sigma_u)</code>	panel-level standard deviation
<code>e(n_quad)</code>	number of quadrature points
<code>e(g_min)</code>	smallest group size
<code>e(g_avg)</code>	average group size
<code>e(g_max)</code>	largest group size
<code>e(p)</code>	<i>p</i> -value for model test
<code>e(rank)</code>	rank of <code>e(V)</code>
<code>e(rank0)</code>	rank of <code>e(V)</code> for constant-only model
<code>e(ic)</code>	number of iterations
<code>e(rc)</code>	return code
<code>e(converged)</code>	1 if converged, 0 otherwise

Macros

<code>e(cmd)</code>	<code>meglm</code>
<code>e(cmd2)</code>	<code>xtlogit</code>
<code>e(cmdline)</code>	command as typed
<code>e(depvar)</code>	name of dependent variable
<code>e(covariates)</code>	list of covariates
<code>e(ivar)</code>	variable denoting groups

e(wtype)	weight type
e(wexp)	weight expression
e(title)	title in estimation output
e(clustvar)	name of cluster variable
e(offset)	linear offset variable
e(chi2type)	Wald or LR; type of model χ^2 test
e(vce)	<i>vcetype</i> specified in <code>vce()</code>
e(vcetype)	title used to label Std. Err.
e(intmethod)	integration method
e(distrib)	Gaussian; the distribution of the random effect
e(opt)	type of optimization
e(which)	max or min; whether optimizer is to perform maximization or minimization
e(ml_method)	type of ml method
e(user)	name of likelihood-evaluator program
e(technique)	maximization technique
e(properties)	b V
e(predict)	program used to implement <code>predict</code>
e(marginsok)	predictions allowed by <code>margins</code>
e(marginswtype)	weight type for <code>margins</code>
e(marginswexp)	weight expression for <code>margins</code>
e(marginsdefault)	default <code>predict()</code> specification for <code>margins</code>
e(marginsdefault)	default <code>predict()</code> specification for <code>margins</code>
e(asbalanced)	factor variables <code>fvset</code> as <code>asbalanced</code>
e(asobserved)	factor variables <code>fvset</code> as <code>asobserved</code>

Matrices

e(b)	coefficient vector
e(Cns)	constraints matrix
e(ilog)	iteration log
e(gradient)	gradient vector
e(cat)	category values
e(V)	variance-covariance matrix of the estimators
e(V_modelbased)	model-based variance

Functions

e(sample)	marks estimation sample
-----------	-------------------------

Methods and formulas

`xtologit` fits via maximum likelihood the random-effects model

$$\Pr(y_{it} > k | \boldsymbol{\kappa}, \mathbf{x}_{it}, \nu_i) = H(\mathbf{x}_{it}\boldsymbol{\beta} + \nu_i - \kappa_k)$$

for $i = 1, \dots, n$ panels, where $t = 1, \dots, n_i$, ν_i are independent and identically distributed $N(0, \sigma_\nu^2)$, and $\boldsymbol{\kappa}$ is a set of cutpoints $\kappa_1, \kappa_2, \dots, \kappa_{K-1}$, where K is the number of possible outcomes; and $H(\cdot)$ is the logistic cumulative distribution function.

From the above, we can derive the probability of observing outcome k for response y_{it} as

$$\begin{aligned} p_{itk} &\equiv \Pr(y_{it} = k | \boldsymbol{\kappa}, \mathbf{x}_{it}, \nu_i) = \Pr(\kappa_{k-1} < \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i + \epsilon_{it} \leq \kappa_k) \\ &= \Pr(\kappa_{k-1} - \mathbf{x}_{it}\boldsymbol{\beta} - \nu_i < \epsilon_{it} \leq \kappa_k - \mathbf{x}_{it}\boldsymbol{\beta} - \nu_i) \\ &= H(\kappa_k - \mathbf{x}_{it}\boldsymbol{\beta} - \nu_i) - H(\kappa_{k-1} - \mathbf{x}_{it}\boldsymbol{\beta} - \nu_i) \\ &= \frac{1}{1 + \exp(-\kappa_k + \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i)} - \frac{1}{1 + \exp(-\kappa_{k-1} + \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i)} \end{aligned}$$

where κ_0 is taken as $-\infty$ and κ_K is taken as $+\infty$. Here \mathbf{x}_{it} does not contain a constant term, because its effect is absorbed into the cutpoints.

We may also express this model in terms of a latent linear response, where observed ordinal responses y_{it} are generated from the latent continuous responses, such that

$$y_{it}^* = \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i + \epsilon_{it}$$

and

$$y_{it} = \begin{cases} 1 & \text{if } y_{it}^* \leq \kappa_1 \\ 2 & \text{if } \kappa_1 < y_{it}^* \leq \kappa_2 \\ \vdots & \\ K & \text{if } \kappa_{K-1} < y_{it}^* \end{cases}$$

The errors ϵ_{it} are distributed as logistic with mean zero and variance $\pi^2/3$ and are independent of ν_i .

Given a set of panel-level random effects ν_i , we can define the conditional distribution for response y_{it} as

$$\begin{aligned} f(y_{it}, \boldsymbol{\kappa}, \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i) &= \prod_{k=1}^K p_{itk}^{I_k(y_{it})} \\ &= \exp \sum_{k=1}^K \left\{ I_k(y_{it}) \log(p_{itk}) \right\} \end{aligned}$$

where

$$I_k(y_{it}) = \begin{cases} 1 & \text{if } y_{it} = k \\ 0 & \text{otherwise} \end{cases}$$

For panel i , $i = 1, \dots, M$, the conditional distribution of $\mathbf{y}_i = (y_{i1}, \dots, y_{in_i})'$ is

$$\prod_{t=1}^{n_i} f(y_{it}, \boldsymbol{\kappa}, \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i)$$

and the panel-level likelihood l_i is given by

$$\begin{aligned} l_i(\boldsymbol{\beta}, \boldsymbol{\kappa}, \sigma_\nu^2) &= \int_{-\infty}^{\infty} \frac{e^{-\nu_i^2/2\sigma_\nu^2}}{\sqrt{2\pi}\sigma_\nu} \left\{ \prod_{t=1}^{n_i} f(y_{it}, \boldsymbol{\kappa}, \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i) \right\} d\nu_i \\ &\equiv \int_{-\infty}^{\infty} g(y_{it}, \boldsymbol{\kappa}, x_{it}, \nu_i) d\nu_i \end{aligned}$$

This integral can be approximated with M -point Gauss–Hermite quadrature

$$\int_{-\infty}^{\infty} e^{-x^2} h(x) dx \approx \sum_{m=1}^M w_m^* h(a_m^*)$$

This is equivalent to

$$\int_{-\infty}^{\infty} f(x) dx \approx \sum_{m=1}^M w_m^* \exp \{ (a_m^*)^2 \} f(a_m^*)$$

where the w_m^* denote the quadrature weights and the a_m^* denote the quadrature abscissas. The log likelihood, L , is the sum of the logs of the panel-level likelihoods l_i .

The default approximation of the log likelihood is by mean–variance adaptive Gauss–Hermite quadrature, which approximates the panel-level likelihood with

$$l_i \approx \sqrt{2}\hat{\sigma}_i \sum_{m=1}^M w_m^* \exp\{(a_m^*)^2\} g(y_{it}, \boldsymbol{\kappa}, x_{it}, \sqrt{2}\hat{\sigma}_i a_m^* + \hat{\mu}_i)$$

where $\hat{\sigma}_i$ and $\hat{\mu}_i$ are the adaptive parameters for panel i . The method of calculating the posterior mean and variance and using those parameters for $\hat{\mu}_i$ and $\hat{\sigma}_i$ is described in detail in [Naylor and Smith \(1982\)](#) and [Skrondal and Rabe-Hesketh \(2004\)](#). We start with $\hat{\sigma}_{i,0} = 1$ and $\hat{\mu}_{i,0} = 0$, and the posterior means and variances are updated in the j th iteration. That is, at the j th iteration of the optimization for l_i , we use

$$l_{i,j} \approx \sum_{m=1}^M \sqrt{2}\hat{\sigma}_{i,j-1} w_m^* \exp\{(a_m^*)^2\} g(y_{it}, \boldsymbol{\kappa}, x_{it}, \sqrt{2}\hat{\sigma}_{i,j-1} a_m^* + \hat{\mu}_{i,j-1})$$

Letting

$$\tau_{i,m,j-1} = \sqrt{2}\hat{\sigma}_{i,j-1} a_m^* + \hat{\mu}_{i,j-1}$$

$$\hat{\mu}_{i,j} = \sum_{m=1}^M (\tau_{i,m,j-1}) \frac{\sqrt{2}\hat{\sigma}_{i,j-1} w_m^* \exp\{(a_m^*)^2\} g(y_{it}, \boldsymbol{\kappa}, x_{it}, \tau_{i,m,j-1})}{l_{i,j}}$$

and

$$\hat{\sigma}_{i,j} = \sum_{m=1}^M (\tau_{i,m,j-1})^2 \frac{\sqrt{2}\hat{\sigma}_{i,j-1} w_m^* \exp\{(a_m^*)^2\} g(y_{it}, \boldsymbol{\kappa}, x_{it}, \tau_{i,m,j-1})}{l_{i,j}} - (\hat{\mu}_{i,j})^2$$

This is repeated until $\hat{\mu}_{i,j}$ and $\hat{\sigma}_{i,j}$ have converged for this iteration of the maximization algorithm. This adaptation is applied on every iteration.

The log likelihood can also be calculated by nonadaptive Gauss–Hermite quadrature with the option `intmethod(ghermite)`, where $\rho = \sigma_\nu^2 / (\sigma_\nu^2 + 1)$:

$$\begin{aligned} L &= \sum_{i=1}^n w_i \log \left\{ \Pr(y_{i1}, \dots, y_{in_i} | \boldsymbol{\kappa}, \mathbf{x}_{i1}, \dots, \mathbf{x}_{in_i}) \right\} \\ &\approx \sum_{i=1}^n w_i \log \left[\frac{1}{\sqrt{\pi}} \sum_{m=1}^M w_m^* \prod_{t=1}^{n_i} f \left\{ y_{it}, \boldsymbol{\kappa}, \mathbf{x}_{it}\boldsymbol{\beta} + a_m^* \left(\frac{2\rho}{1-\rho} \right)^{1/2} \right\} \right] \end{aligned}$$

Both quadrature formulas require that the integrated function be well approximated by a polynomial of degree equal to the number of quadrature points. The number of periods (panel size) can affect whether

$$\prod_{t=1}^{n_i} f(y_{it}, \boldsymbol{\kappa}, \mathbf{x}_{it}\boldsymbol{\beta} + \nu_i)$$

is well approximated by a polynomial. As panel size and ρ increase, the quadrature approximation can become less accurate. For large ρ , the random-effects model can also become unidentified. Adaptive quadrature gives better results for correlated data and large panels than nonadaptive quadrature; however, we recommend that you use the `quadchk` command (see [\[XT\] quadchk](#)) to verify the quadrature approximation used in this command, whichever approximation you choose.

xtologit and the robust VCE estimator

Specifying `vce(robust)` or `vce(cluster clustvar)` causes the Huber/White/sandwich VCE estimator to be calculated for the coefficients estimated in this regression. See [P] [_robust](#), particularly [Introduction](#) and [Methods and formulas](#). Wooldridge (2016) and Arellano (2003) discuss this application of the Huber/White/sandwich VCE estimator. As discussed by Wooldridge (2016), Stock and Watson (2008), and Arellano (2003), specifying `vce(robust)` is equivalent to specifying `vce(cluster panelvar)`, where *panelvar* is the variable that identifies the panels.

Clustering on the panel variable produces a consistent VCE estimator when the disturbances are not identically distributed over the panels or there is serial correlation in ϵ_{it} .

The cluster-robust VCE estimator requires that there are many clusters and the disturbances are uncorrelated across the clusters. The panel variable must be nested within the cluster variable because of the within-panel correlation that is generally induced by the random-effects transform when there is heteroskedasticity or within-panel serial correlation in the idiosyncratic errors.

References

- Allison, P. D. 2009. *Fixed Effects Regression Models*. Newbury Park, CA: Sage.
- Arellano, M. 2003. *Panel Data Econometrics*. Oxford: Oxford University Press.
- Conway, M. R. 1990. A random effects model for binary data. *Biometrics* 46: 317–328.
- Flay, B. R., B. R. Brannon, C. A. Johnson, W. B. Hansen, A. L. Ulene, D. A. Whitney-Saltiel, L. R. Gleason, S. Sussman, M. D. Gavin, K. M. Glowacz, D. F. Sobol, and D. C. Spiegel. 1988. The television, school, and family smoking cessation and prevention project: I. Theoretical basis and program development. *Preventive Medicine* 17: 585–607.
- Liang, K.-Y., and S. L. Zeger. 1986. Longitudinal data analysis using generalized linear models. *Biometrika* 73: 13–22.
- Naylor, J. C., and A. F. M. Smith. 1982. Applications of a method for the efficient computation of posterior distributions. *Journal of the Royal Statistical Society, Series C* 31: 214–225.
- Neuhaus, J. M. 1992. Statistical methods for longitudinal and clustered designs with binary responses. *Statistical Methods in Medical Research* 1: 249–273.
- Neuhaus, J. M., J. D. Kalbfleisch, and W. W. Hauck. 1991. A comparison of cluster-specific and population-averaged approaches for analyzing correlated binary data. *International Statistical Review* 59: 25–35.
- Pendergast, J. F., S. J. Gange, M. A. Newton, M. J. Lindstrom, M. Palta, and M. R. Fisher. 1996. A survey of methods for analyzing clustered binary response data. *International Statistical Review* 64: 89–118.
- Rabe-Hesketh, S., and A. Skrondal. 2012. *Multilevel and Longitudinal Modeling Using Stata*. 3rd ed. College Station, TX: Stata Press.
- Skrondal, A., and S. Rabe-Hesketh. 2004. *Generalized Latent Variable Modeling: Multilevel, Longitudinal, and Structural Equation Models*. Boca Raton, FL: Chapman & Hall/CRC.
- Stock, J. H., and M. W. Watson. 2008. Heteroskedasticity-robust standard errors for fixed effects panel data regression. *Econometrica* 76: 155–174.
- Twisk, J. W. R. 2013. *Applied Longitudinal Data Analysis for Epidemiology: A Practical Guide*. 2nd ed. Cambridge: Cambridge University Press.
- Wooldridge, J. M. 2016. *Introductory Econometrics: A Modern Approach*. 6th ed. Boston: Cengage.

Also see

- [XT] [xtologit postestimation](#) — Postestimation tools for xtologit
- [XT] [quadchk](#) — Check sensitivity of quadrature approximation
- [XT] [xtoprobit](#) — Random-effects ordered probit models
- [XT] [xtset](#) — Declare data to be panel data
- [ME] [meologit](#) — Multilevel mixed-effects ordered logistic regression
- [R] [logistic](#) — Logistic regression, reporting odds ratios
- [R] [logit](#) — Logistic regression, reporting coefficients
- [U] [20 Estimation and postestimation commands](#)