

SCOREDUM - A stata command to test for fixed effects in Poisson and Logit models.

Paulo Guimaraes

CEFUP - Universidade do Porto

September 2012

Maximum Likelihood Estimation

- Suppose we have a probability model $f(Y; \mathbf{X}, \boldsymbol{\theta})$
- Y is the dependent variable, $\boldsymbol{\theta}$ is a vector of k unknown parameters and \mathbf{X} is a set of covariates.
- Let $\mu_i = \mathbf{x}_i' \boldsymbol{\theta}$
- The likelihood function is:

$$L(\boldsymbol{\theta}; \mathbf{Y}, \mathbf{X}) = \prod_{i=1}^N f(\boldsymbol{\theta}; y_i, \mathbf{x}_i)$$

- The m.l.e is the value $\hat{\boldsymbol{\theta}}$ that maximizes $L(\boldsymbol{\theta}; \mathbf{Y}, \mathbf{X})$

Testing Nested Hypothesis

- Suppose we want to test the null hypothesis $\mathbf{h}(\boldsymbol{\theta}) = 0$
- There are 3 likelihood based procedures for testing nested hypothesis:
 - Likelihood Ratio Tests
 - Wald Tests
 - Score Tests (or Lagrange Multiplier)
- These tests are all asymptotically chi-squared distributed with degrees of freedom equal to the number of restrictions imposed
- Let $\hat{\boldsymbol{\theta}}_r$ be the maximum likelihood estimator under the null hypothesis (the restricted MLE) and $\hat{\boldsymbol{\theta}}_u$ under the alternative (the unrestricted MLE).

- The Likelihood Ratio Test

$$LR = -2 \left[\ln L(\hat{\theta}_r) - \ln L(\hat{\theta}_u) \right]$$

- The Wald Test

$$W = \mathbf{h}(\hat{\theta}_u)' \left[V \left[\mathbf{h}(\hat{\theta}_u) \right] \right]^{-1} \mathbf{h}(\hat{\theta}_u)$$

- The Score Test

$$LM = \mathbf{s}(\hat{\theta}_r)' \left[\mathbf{I}(\hat{\theta}_r) \right]^{-1} \mathbf{s}(\hat{\theta}_r)$$

Testing Group Comparisons

- Suppose the all N observations are classified into G mutually exclusive groups
- Groups can be panels or clusters
- Let $\theta' = [\alpha', \beta']$
- α is a subset of θ with k_1 elements.
- $k = k_1 + k_2$ elements
- We want to test the following hypothesis

$$H_0 : \alpha_1 = \alpha_2 = \dots = \alpha_G .$$

The Score Test for Group Comparisons

- The score test is:

$$T = \mathbf{s}(\hat{\boldsymbol{\vartheta}})' \left[-\mathbf{H}(\hat{\boldsymbol{\vartheta}}) \right]^{-1} \mathbf{s}(\hat{\boldsymbol{\vartheta}})$$

the score equations and Hessian are calculated with respect to all the coefficients in $\boldsymbol{\vartheta}' = [\boldsymbol{\alpha}'_1, \boldsymbol{\alpha}'_2, \dots, \boldsymbol{\alpha}'_G; \boldsymbol{\beta}']$, but evaluated at

$$\hat{\boldsymbol{\vartheta}}' = [\hat{\boldsymbol{\alpha}}', \hat{\boldsymbol{\alpha}}', \dots, \hat{\boldsymbol{\alpha}}', \hat{\boldsymbol{\beta}}']$$

- The $\hat{\boldsymbol{\vartheta}}'$ are the m.l.e. solutions obtained under the null hypothesis
- The score test is asymptotically distributed as chi-square with $k_1(G - 1)$ degrees of freedom

The Score Test for Group Comparisons

- Partitioning the score and Hessian with respect to α and β we get

$$T = - \begin{bmatrix} \mathbf{s}_\alpha \\ \mathbf{s}_\beta \end{bmatrix}' \begin{bmatrix} \mathbf{H}_{\alpha\alpha} & \mathbf{H}_{\alpha\beta} \\ \mathbf{H}_{\beta\alpha} & \mathbf{H}_{\beta\beta} \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{s}_\alpha \\ \mathbf{s}_\beta \end{bmatrix}$$

- Since $\mathbf{s}_\beta(\hat{\vartheta}) = \mathbf{0}$ the score test may be written as:

$$T = -\mathbf{s}'_\alpha \left[\mathbf{H}_{\alpha\alpha} + \mathbf{H}_{\alpha\beta} \left[-\mathbf{H}_{\beta\beta} \right]^{-1} \mathbf{H}_{\beta\alpha} \right]^{-1} \mathbf{s}_\alpha$$

$$T = -\mathbf{s}'_\alpha \left[\mathbf{H}_{\alpha\alpha}^{-1} + \mathbf{H}_{\alpha\alpha}^{-1} \mathbf{H}_{\alpha\beta} \left[\mathbf{H}_{\beta\beta} - \mathbf{H}_{\beta\alpha} \mathbf{H}_{\alpha\alpha}^{-1} \mathbf{H}_{\alpha\beta} \right]^{-1} \mathbf{H}_{\beta\alpha} \mathbf{H}_{\alpha\alpha}^{-1} \right] \mathbf{s}_\alpha$$

- \mathbf{s}_α , $\mathbf{H}_{\alpha\alpha}$, $\mathbf{H}_{\alpha\beta}$ and $\mathbf{H}_{\beta\beta}$ can be easily calculated from the restricted model!

- If $k_2 = 0$ or $G = N$ then

$$T = -\mathbf{s}'_{\alpha} \mathbf{H}_{\alpha\alpha}^{-1} \mathbf{s}_{\alpha}$$

- Pearson tests are particular cases

$$T = - \sum_{g=1}^G \frac{s_{\alpha, \bullet g}^2}{h_{\alpha\alpha, \bullet g}}$$

- If Y is Poisson: $T = \sum_{g=1}^G \frac{(y_{\bullet g} - n_g \bar{y})^2}{n_g \bar{y}}$
- Y is Bernoulli: $T = \sum_{g=1}^G \frac{(y_{\bullet g} - n_g \bar{p})^2}{n_g \bar{p}(1-\bar{p})}$
- If Y is normal: $T = \sum_{g=1}^G \frac{n_g (\bar{y}_g - \bar{y})^2}{\sigma^2}$

Syntax

```
scoredum [indepvars], group(varname) [options]
```

Examples:

- Test for fixed effects in Poisson Regression

```
poisson y x1 x2 x3  
scoredum, group(grvar)
```
- Test for "slippery slope" in x1

```
scoredum x1, group(grvar)
```
- Test for differences in several coefficients across groups

```
scoredum x1 x2 x3, group(grvar) cons
```
- Pearson chi-square test for count data

```
poisson y  
scoredum, group(grvar)
```